

Q1. The current voltage relation of diode is given by $I = (e^{1000 V/T} - 1) \text{ mA}$, where the applied voltage V is in volts and the temperature T is in degree Kelvin. If a student makes an error measuring $\pm 0.01 \text{ V}$ while measuring the current of 5 mA at 300 K , what will be the error in the value of current in mA ?

- (1) 0.2 mA (2) 0.02 mA
 (3) 0.5 mA (4) 0.05 mA

Q2. From a tower of height H , a particle is thrown vertically upwards with a speed u . The time taken by the particle, to hit the ground, is n times that taken by it to reach the highest point of its path. The relation between H , u and n is :

- (1) $2gH = n^2u^2$ (2) $gH = (n-2)^2u^2$
 (3) $2gH = nu^2(n-2)$ (4) $gH = (n-2)u^2$

Q3. A block of mass m is placed on a surface with a vertical cross section given by $y = \frac{x^3}{6}$. If the coefficient of friction is 0.5 , the maximum height above the ground at which the block can be placed without slipping is


- (1) $\frac{1}{6} \text{ m}$ (2) $\frac{2}{3} \text{ m}$
 (3) $\frac{1}{3} \text{ m}$ (4) $\frac{1}{2} \text{ m}$

Q4. When a rubber-band is stretched by a distance x , it exerts a restoring force of magnitude

$F = ax + bx^2$ where a and b are constants. The work done in stretching the unstretched rubber-band by L is :

- (1) $aL^2 + bL^3$ (2) $\frac{1}{2}(aL^2 + bL^3)$
 (3) $\frac{aL^2}{2} + \frac{bL^3}{3}$ (4) $\frac{1}{2}\left(\frac{aL^2}{2} + \frac{bL^3}{3}\right)$

Q5. A mass m is supported by a massless string wound around a uniform hollow cylinder of mass m and radius R . If the string does not slip on the cylinder, with what acceleration will the mass fall on release?



- (1) $\frac{2g}{3}$ (2) $\frac{g}{2}$
 (3) $\frac{5g}{6}$ (4) g

Q6. A bob of mass m attached to an inextensible string of length l is suspended from a vertical support. The bob rotates in a horizontal circle with an angular speed $\omega \text{ rad/s}$ about the vertical. About the point of suspension :

- (1) Angular momentum is conserved (2) Angular momentum changes in magnitude but not in direction
 (3) Angular momentum changes in direction but not in magnitude (4) Angular momentum changes both in direction and magnitude

Q7. Four particles, each of mass M and equidistant from each other, move along a circle of radius R under the action of their mutual gravitational attraction. The speed of each particle is

(1) $\sqrt{\frac{GM}{R}}$


(2) $\sqrt{2\sqrt{2}\frac{GM}{R}}$

(3) $\sqrt{\frac{GM}{R}(1+2\sqrt{2})}$

(4) $\frac{1}{2}\sqrt{\frac{GM}{R}(1+2\sqrt{2})}$

Q8. There is a circular tube in a vertical plane. Two liquids which do not mix and of densities d_1 and d_2 are filled in the tube. Each liquid subtends 90° angle at centre. Radius joining their interface makes an angle α with vertical.

Ratio $\frac{d_1}{d_2}$ is :



(1) $\frac{1+\sin\alpha}{1-\sin\alpha}$


(2) $\frac{1+\cos\alpha}{1-\cos\alpha}$

(3) $\frac{1+\tan\alpha}{1-\tan\alpha}$

(4) $\frac{1+\sin\alpha}{1-\cos\alpha}$

Q9. On heating water, bubbles being formed at the bottom of the vessel detach and rise. Take the bubbles to be spheres of radius R and making a circular contact of radius r with the bottom of the vessel. If $r \ll R$, and the surface tension of water is T , value of r just before bubbles detach is :

(density of water is ρ_w)



(1) $R^2 \sqrt{\frac{2\rho_w g}{3T}}$

(2) $R^2 \sqrt{\frac{\rho_w g}{6T}}$

(3) $R^2 \sqrt{\frac{\rho_w g}{T}}$

(4) $R^2 \sqrt{\frac{3\rho_w g}{T}}$

Q10. An open glass tube is immersed in mercury in such a way that a length of 8 cm extends above the mercury level. The open end of the tube is then closed and sealed and the tube is raised vertically up by additional 46 cm. What will be length of the air column above mercury in the tube now ?

(Atmospheric pressure = 76 cm of Hg)

(1) 16 cm

(2) 22 cm

(3) 38 cm

(4) 6 cm

Q11. The pressure that has to be applied to the ends of a steel wire of length 10 cm to keep its length constant when its temperature is raised by 100°C is :

(For steel, Young's modulus is $2 \times 10^{11} \text{ N m}^{-2}$ and coefficient of thermal expansion is $1.1 \times 10^{-5} \text{ K}^{-1}$)

(1) $2.2 \times 10^8 \text{ Pa}$

(2) $2.2 \times 10^9 \text{ Pa}$

(3) $2.2 \times 10^7 \text{ Pa}$


(4) $2.2 \times 10^6 \text{ Pa}$

Q12. Three rods of Copper, Brass and Steel are welded together to form a Y-shaped structure. Area of cross-section of each rod is 4 cm^2 . End of copper rod is maintained at 100°C . Where as ends of brass and steel are kept at

0°C . Lengths of the copper, brass and steel rods are 46, 13 and 12 cms respectively. The rods are thermally insulated from surroundings except at ends. Thermal conductivities of copper, brass and steel are 0.92, 0.26 and 0.12 CGS units respectively. Rate of heat flow through copper rod is :

- (1) 1.2 Cal/s (2) 2.4 Cal/s
(3) 4.8 Cal/s (4) 6.0 Cal/s

Q13. One mole of diatomic ideal gas undergoes a cyclic process ABC as shown in figure. The process BC is adiabatic. The temperatures at A, B and C are 400 K, 800 K and 600 K respectively. Choose the correct statement :



- (1) The change in internal energy in whole cyclic process is 250 R
(2) The change in internal energy in the process CA is 700 R.
(3) The change in internal energy in the process AB is -350 R
(4) The change in internal energy in the process BC is -500 R

Q14. A particle moves with simple harmonic motion in a straight line. In first τ s, after starting from rest it travels a distance a , and in next τ s it travels $2a$, in same direction, then :

- (1) Amplitude of motion is $3a$ (2) Time period of oscillations is 8τ
(3) Amplitude of motion is $4a$ (4) Time period of oscillations is 6τ

Q15. A pipe of length 85 cm is closed from one end. Find the number of possible natural oscillations of air column in the pipe whose frequencies lie below 1250 Hz. The velocity of sound in air is 340 m/s.

- (1) 12 (2) 8
(3) 6 (4) 4

Q16. Assume that an electric field $\vec{E} = 30x^2\hat{i}$ exists in space. Then the potential difference $V_A - V_O$, where V_O is the potential at the origin and V_A the potential at $x = 2$ m is :

- (1) 120 J C^{-1} (2) -120 J C^{-1}
(3) -80 J C^{-1} (4) 80 J C^{-1}

Q17. A parallel plate capacitor is made of two circular plates separated by a distance of 5 mm and with a dielectric of dielectric constant 2.2 between them. When the electric field in the dielectric is $3 \times 10^4 \text{ V/m}$, the charge density of the positive plate will be close to :

- (1) $6 \times 10^{-7} \text{ C/m}^2$ (2) $3 \times 10^{-7} \text{ C/m}^2$
(3) $3 \times 10^4 \text{ C/m}^2$ (4) $6 \times 10^4 \text{ C/m}^2$


Q18. In a large building, there are 15 bulbs of 40 W, 5 bulbs of 100 W, 5 fans of 80 W and 1 heater of 1 kW. The voltage of the electric mains is 220 V. The minimum capacity of the main fuse of the building will be:

- (1) 8 A (2) 10 A
(3) 12 A (4) 14 A

Q19. The coercivity of a small magnet, where the ferromagnet gets demagnetised is 3×10^3 A/m. The current required to be passed in a solenoid of length 10 cm and number of turns 100, so that the magnet gets demagnetised when inside the solenoid is


- (1) 30 mA (2) 60 mA
(3) 3 A (4) 6 A

Q20. A conductor lies along the z-axis at $-1.5 \leq z \leq 1.5$ m and carries a fixed current of 10.0 A in $-\hat{a}_z$ direction (see figure). For a field $\vec{B} = 3.0 \times 10^{-4} e^{-0.2x} \hat{a}_y$ T, find the power required to move the conductor at constant speed to $x = 2.0$ m, $y = 0$ m in 5×10^{-3} s. Assume parallel motion along the x-axis.



- (1) 1.57 W (2) 2.97 W
(3) 14.85 W (4) 29.7 W

Q21. In the circuit shown here, the point 'C' is kept connected to point 'A' till the current flowing through the circuit becomes constant. Afterward, suddenly, point 'C' is disconnected from point 'A' and connected to point 'B' at time $t = 0$. Ratio of the voltage across resistance and the inductor at $t = \frac{L}{R}$ will be equal to :



- (1) $\frac{e}{1-e}$ (2) 1
(3) -1 (4) $\frac{1-e}{e}$

Q22. During the propagation of electromagnetic wave in a particular medium :-

- (1) Electric energy density is double of the magnetic energy density (2) Electric energy density is half of the magnetic energy density
(3) Electric energy density is equal to the magnetic energy density (4) Both electric and magnetic energy densities are zero

Q23. Match List - I (Electromagnetic wave type) with List - II (Its association/application) and select the correct option from the choices given below the lists :

List - I

List - II

- | | |
|----------------------|--|
| (a) Infrared waves | (i) To treat muscular strain |
| (b) Radio waves | (ii) For broadcasting |
| (c) X - rays | (iii) To detect fracture of bones |
| (d) Ultraviolet rays | (iv) Absorbed by the ozone layer of the atmosphere |

(1) (a) (b) (c) (d)

(2) (a) (b) (c) (d)

(iv) (iii) (ii) (i)

(i) (ii) (iv) (iii)

(3) (a) (b) (c) (d)

(4) (a) (b) (c) (d)

(iii) (ii) (i) (iv)

(i) (ii) (iii) (iv)

Q24. A thin convex lens made from crown glass ($\mu = \frac{3}{2}$) has focal length f . When it is measured in two different liquids having refractive indices $\frac{4}{3}$ and $\frac{5}{3}$, it has the focal lengths f_1 and f_2 respectively. The correct relation between the focal lengths is :

(1) $f_1 = f_2 < f$ (2) $f_1 > f$ and f_2 becomes negative(3) $f_2 > f$ and f_1 becomes negative(4) f_1 and f_2 both become negative

Q25. A green light is incident from the water to the air - water interface at the critical angle (θ_c). Select the correct statement.

(1) The entire spectrum of visible light will come out of the water at an angle of 90° to the normal

(2) The spectrum of visible light whose frequency is less than that of green light will come out to the air medium

(3) The spectrum of visible light whose frequency is more than that of green light will come out to the air medium

(4) The entire spectrum of visible light will come out of the water at various angles to the normal

Q26. Two beams, A and B of plane polarized light with mutually perpendicular planes of polarization are seen through a polaroid. From the position when the beam A has maximum intensity (and beam B has zero intensity), a rotation of polaroid through 30° makes the two beams appear equally bright. If the initial intensities of the two beams are I_A and I_B respectively, then $\frac{I_A}{I_B}$ equals :

(1) 3

(2) $\frac{3}{2}$

(3) 1

(4) $\frac{1}{3}$

Q27. The radiation corresponding to $3 \rightarrow 2$ transition of hydrogen atom falls on a metal surface to produce photoelectrons. These electrons are made to enter a magnetic field of 3×10^{-4} T. If the radius of the largest circular path followed by these electrons is 10.0 mm, the work function of the metal is close to :

(1) 1.8 eV

(2) 1.1 eV



(3) 0.8 eV

(4) 1.6 eV

Q28. Hydrogen (${}_1\text{H}^1$), Deuterium (${}_1\text{H}^2$), singly ionised Helium (${}_2\text{He}^4$)⁺ and doubly ionised lithium (${}_3\text{Li}^6$)⁺⁺ all have one electron around the nucleus. Consider an electron transition from $n = 2$ to $n = 1$. If the wave lengths of emitted radiation are λ_1 , λ_2 , λ_3 and λ_4 respectively then approximately which one of the following is correct ?

(1) $4\lambda_1 = 2\lambda_2 = 2\lambda_3 = \lambda_4$ (2) $\lambda_1 = 2\lambda_2 = 2\lambda_3 = \lambda_4$ (3) $\lambda_1 = \lambda_2 = 4\lambda_3 = 9\lambda_4$ (4) $\lambda_1 = 2\lambda_2 = 3\lambda_3 = 4\lambda_4$

Q29. The forward biased diode connection is :



Q30. A student measured the length of a rod and wrote it as 3.50 cm. Which instrument did he use to measure it ?

(1) A meter scale

(2) A vernier calliper where the 10 divisions in vernier scale matches with 9 division in main scale and main scale has 10 divisions in 1 cm

(3) A screw gauge having 100 divisions in the circular scale and pitch as 1 mm

(4) A screw gauge having 50 divisions in the circular scale and pitch as 1 mm

Q31. The ratio of masses of oxygen and nitrogen in a particular gaseous mixture is 1 : 4. The ratio of number of their molecules is:

(1) 1 : 4

(2) 7 : 32

(3) 1 : 8

(4) 3 : 16

Q32. The correct set of four quantum numbers for the valence electrons of rubidium atom ($Z = 37$) is

(1) 5, 0, 0, $+\frac{1}{2}$

(2) 5, 1, 0, $+\frac{1}{2}$

(3) 5, 1, 1, $+\frac{1}{2}$

(4) 5, 0, 1, $+\frac{1}{2}$

Q33. Which one of the following properties is **not** shown by NO ?





(1) It is diamagnetic in gaseous state

(2) It is a neutral oxide

(3) It combines with oxygen to form nitrogen dioxide

(4) It's bond order is 2.5

Q34. For which of the following molecule significant $\mu \neq 0$?



(1) Only (a)

(2) (a) and (b)

(3) Only (c)

(4) (c) and (d)

Q35. If Z is the compressibility factor, then Van der Waal's equation at low pressure can be written as:


$$(1) Z = 1 + \frac{RT}{P_b}$$

$$(3) Z = 1 - \frac{P_b}{RT}$$

$$(2) Z = 1 - \frac{a}{VRT}$$

$$(4) Z = 1 + \frac{P_b}{RT}$$

Q36. For complete combustion of ethanol,



the amount of heat produced as measured in bomb calorimeter, is $1364.47 \text{ kJ mol}^{-1}$ at 25°C . Assuming ideality the Enthalpy of combustion, $\Delta_c H$, for the reaction will be: ($R = 8.314 \text{ kJ mol}^{-1}$)

$$(1) -1366.95 \text{ kJ mol}^{-1}$$

$$(3) -1460.50 \text{ kJ mol}^{-1}$$

$$(2) -1361.95 \text{ kJ mol}^{-1}$$

$$(4) -1350.50 \text{ kJ mol}^{-1}$$

Q37. For the reaction $SO_2(g) + \frac{1}{2}O_2(g) \rightleftharpoons SO_3(g)$, if $K_P = K_C(RT)^x$ where the symbols have usual meaning then the value of x is:
(assuming ideality)




$$(1) -1$$

$$(3) \frac{1}{2}$$

$$(2) -\frac{1}{2}$$

$$(4) 1$$

Q38. In which of the following reactions H_2O_2 acts as a reducing agent ?



(1) (a), (b)

(2) (c), (d)

(3) (a), (c)

(4) (b), (d)

Q39. For the estimation of nitrogen, 1.4 g of an organic compound was digested by the Kjeldahl method and the evolved ammonia was absorbed in 60 ml of $\frac{M}{10}$ sulphuric acid. The unreacted acid required 20 ml of $\frac{M}{10}$ sodium hydroxide for complete neutralization. The percentage of nitrogen in the compound is

$$(1) 6\%$$

$$(3) 3\%$$

$$(2) 10\%$$

$$(4) 5\%$$

Q40. Considering the basic strength of amines in an aqueous solution, which one has the smallest pK_b value?



Q41. The major organic compound formed by the reaction of 1, 1, 1 - trichloroethane with silver powder is :

(1) Acetylene

(2) Ethene

(3) 2 - Butyne

(4) 2 - Butene

Q42. CsCl crystallises in body centred cubic lattice. If 'a' is its edge length then which of the following expressions is correct?

$$(1) r_{Cs^+} + r_{Cl^-} = 3a$$

$$(2) r_{Cs^+} + r_{Cl^-} = \frac{3a}{2}$$

$$(3) r_{Cs^+} + r_{Cl^-} = \frac{\sqrt{3}}{2} a$$

$$(4) r_{Cs^+} + r_{Cl^-} = \sqrt{3}a$$

Q43. Consider separate solutions of 0.500 M C_2H_5OH (aq), 0.100 M $Mg_3(PO_4)_2$ (aq), 0.250 M KBr (aq) and 0.125 M Na_3PO_4 (aq) at 25°C . Which statement is true about these solutions, assuming all salts to be

strong electrolytes?

- (1) They all have the same osmotic pressure. (2) 0.100 M $\text{Mg}_3(\text{PO}_4)_2$ (aq) has the highest osmotic pressure.
 (3) 0.125 M Na_3PO_4 (aq) has the highest osmotic pressure. (4) 0.500 M $\text{C}_2\text{H}_5\text{OH}$ (aq) has the highest osmotic pressure.

Q44. Resistance of 0.2 M solution of an electrolyte is $50\ \Omega$. The specific conductance of the solution is $1.4\ \text{S m}^{-1}$.

The resistance of 0.5 M solution of the same electrolyte is $280\ \Omega$. The molar conductivity of 0.5 M solution of the electrolyte in $\text{S m}^2\ \text{mol}^{-1}$ is :

- (1) 5×10^{-4} (2) 5×10^{-3}
 (3) 5×10^3 (4) 5×10^2

Q45. The equivalent conductance of NaCl at concentration C and at infinite dilution are λ_C and λ_∞ , respectively.

The correct relationship between λ_C and λ_∞ is given as :



(where the constant B is positive)

- (1) $\lambda_C = \lambda_\infty + (B)C$ (2) $\lambda_C = \lambda_\infty - (B)C$
 (3) $\lambda_C = \lambda_\infty - (B)\sqrt{C}$ (4) $\lambda_C = \lambda_\infty + (B)\sqrt{C}$

Q46. The metal that cannot be obtained by the electrolysis of an aqueous solution of its salt is

- (1) Ag (2) Ca
 (3) Cu (4) Cr

Q47. Given below are the half - cell reactions :



The E° for $3\text{Mn}^{2+} \rightarrow \text{Mn} + 2\text{Mn}^{3+}$ will be :

- (1) $-2.69\ \text{V}$; the reaction will not occur (2) $-2.69\ \text{V}$; the reaction will occur
 (3) $-0.33\ \text{V}$; the reaction will not occur (4) $-0.33\ \text{V}$; the reaction will occur

Q48. For the non-stoichiometry reaction, $2\text{A} + \text{B} \rightarrow \text{C} + \text{D}$, the following kinetic data were obtained in three separate experiments, all at 298 K.

Initial Concentration Initial Concentration Initial rate of formation of C

(A) (B) (mol L⁻¹ S⁻¹)

0.1 M 0.1 M 1.2×10^{-3}


0.1 M 0.2 M 1.2×10^{-3}

0.2 M 0.1 M 2.4×10^{-3}

The rate law for the formation of C is

- (1) $\frac{dc}{dt} = k[\text{A}][\text{B}]$ (2) $\frac{dc}{dt} = k[\text{A}]^2[\text{B}]$
 (3) $\frac{dc}{dt} = k[\text{A}][\text{B}]^2$ (4) $\frac{dc}{dt} = k[\text{A}]$

Q49. Which series of reactions correctly represents the chemical relations related to iron and its compound?



Q50. Among the following oxoacids, the correct decreasing order of acid strength is :

- (1) $\text{HOCl} > \text{HClO}_2 > \text{HClO}_3 > \text{HClO}_4$ (2) $\text{HClO}_4 > \text{HOCl} > \text{HClO}_2 > \text{HClO}_3$
 (3) $\text{HClO}_4 > \text{HClO}_3 > \text{HClO}_2 > \text{HOCl}$ (4) $\text{HClO}_2 > \text{HClO}_4 > \text{HClO}_3 > \text{HOCl}$

Q51. The equation which is balanced and represents the correct product(s) is

- (1) $\text{Li}_2\text{O} + 2\text{KCl} \rightarrow 2\text{LiCl} + \text{K}_2\text{O}$ (2) $[\text{CoCl}(\text{NH}_3)_5]^+ + 5\text{H}^+ \rightarrow \text{Co}^{2+} + 5\text{NH}_4^+ + \text{Cl}^-$
 (3) $[\text{Mg}(\text{H}_2\text{O})_6]^{2+} + (\text{EDTA})^{4-} \xrightarrow{\text{excess NaOH}} [\text{Mg}(\text{EDTA})]^{2-} + 6\text{H}_2\text{O}$ (4) $\text{CuSO}_4 + 4\text{KCN} \rightarrow \text{K}_2[\text{Cu}(\text{CN})_4] + \text{K}_2\text{SO}_4$

Q52. The octahedral complex of a metal ion M^{3+} with four monodentate ligands L_1 , L_2 , L_3 and L_4 absorb wavelengths in the region of red, green, yellow and blue, respectively. The increasing order of ligand strength of the four ligands is

- (1) $\text{L}_4 < \text{L}_3 < \text{L}_2 < \text{L}_1$ (2) $\text{L}_1 < \text{L}_3 < \text{L}_2 < \text{L}_4$
 (3) $\text{L}_3 < \text{L}_2 < \text{L}_4 < \text{L}_1$ (4) $\text{L}_1 < \text{L}_2 < \text{L}_4 < \text{L}_3$

Q53. The correct statement for the molecule CsI_3 , is :

- (1) It is a covalent molecule. (2) It contains Cs^+ and I_3^- ions.
 (3) It contains Cs^{3+} and I^- ions. (4) It contains Cs^+ , I^- and lattice I_2 molecule.


Q54. In $\text{S}_{\text{N}}2$ reactions, the correct order of reactivity for the following compounds :





- CH_3Cl , $\text{CH}_3\text{CH}_2\text{Cl}$, $(\text{CH}_3)_2\text{CHCl}$ and $(\text{CH}_3)_3\text{CCl}$ is :
 (1) $\text{CH}_3\text{Cl} > (\text{CH}_3)_2\text{CHCl} > \text{CH}_3\text{CH}_2\text{Cl} > (\text{CH}_3)_3\text{CCl}$ (2) $\text{CH}_3\text{Cl} > \text{CH}_3\text{CH}_2\text{Cl} > (\text{CH}_3)_2\text{CHCl} > (\text{CH}_3)_3\text{CCl}$
 (3) $\text{CH}_3\text{CH}_2\text{Cl} > \text{CH}_3\text{Cl} > (\text{CH}_3)_2\text{CHCl} > (\text{CH}_3)_3\text{CCl}$ (4) $(\text{CH}_3)_2\text{CHCl} > \text{CH}_3\text{CH}_2\text{Cl} > \text{CH}_3\text{Cl} > (\text{CH}_3)_3\text{CCl}$

Q55. The most suitable reagent for the conversion of $\text{R}-\text{CH}_2-\text{OH} \rightarrow \text{R}-\text{CHO}$ is :


- (1) KMnO_4 (2) $\text{K}_2\text{Cr}_2\text{O}_7$
 (3) CrO_3 (4) PCC (Pyridinium Chlorochromate)

Q56. What is the product "C" after following reactions -





Q57. In the reaction,



- (1) Acetaldehyde (2) Acetylene
(3) Ethylene (4) Acetyl chloride

Q58. On heating an aliphatic primary amine with chloroform and ethanolic potassium hydroxide, the organic compound formed is :

- (1) An alkanol (2) An alkanediol
(3) An alkyl cyanide (4) An alkyl isocyanide

Q59. Which among the following is classified as a condensation polymer?

- (1) Dacron (2) Neoprene
(3) Teflon (4) Acrylonitrile

Q60. Which one of the following bases is **not** present in DNA ?

- (1) Quinoline (2) Adenine
(3) Cytosine (4) Thymine

Q61. If $a \in R$ and the equation $-3(x - [x])^2 + 2(x - [x]) + a^2 = 0$ (where $[x]$ denotes the greatest integer $\leq x$) has no integral solution, then all possible values of a lie in the interval

- (1) $(-2, -1)$ (2) $(-\infty, -2) \cup (2, \infty)$
(3) $(-1, 0) \cup (0, 1)$ (4) $(1, 2)$

Q62. Let α and β be the roots of equation $px^2 + qx + r = 0$, $p \neq 0$. If p, q, r are in A.P. and $\frac{1}{\alpha} + \frac{1}{\beta} = 4$, then the value of $|\alpha - \beta|$ is

- (1) $\frac{\sqrt{34}}{9}$ (2) $\frac{2\sqrt{13}}{9}$
(3) $\frac{\sqrt{61}}{9}$ (4) $\frac{2\sqrt{17}}{9}$

Q63. If z is a complex number such that $|z| \geq 2$, then the minimum value of $|z + \frac{1}{z}|$:

- (1) Is strictly greater than $\frac{5}{2}$ (2) Is strictly greater than $\frac{3}{2}$ but less than $\frac{5}{2}$
(3) Is equal to $\frac{5}{2}$ (4) Lies in the interval $(1, 2)$

Q64. If $(10)^9 + 2(11)^1(10)^8 + 3(11)^2(10)^7 + \dots + 10(11)^9 = k(10)^9$, then k is equal to :

(1) 100

(3) $\frac{121}{10}$

(2) 110

(4) $\frac{441}{100}$

Q65. Three positive numbers form an increasing $G. P.$ If the middle term in this $G. P.$ is doubled, the new numbers are in $A. P.$ Then the common ratio of the $G. P.$ is :

(1) $2 - \sqrt{3}$

(3) $\sqrt{2} + \sqrt{3}$

(2) $2 + \sqrt{3}$

(4) $3 + \sqrt{2}$

Q66. If the coefficients of x^3 and x^4 in the expansion of $(1 + ax + bx^2)(1 - 2x)^{18}$ in powers of x are both zero, then (a, b) is equal to

(1) $(14, \frac{272}{3})$

(3) $(16, \frac{251}{3})$

(2) $(16, \frac{272}{3})$

(4) $(14, \frac{251}{3})$

Q67. Let $f_k(x) = \frac{1}{k}(\sin^k x + \cos^k x)$ where $x \in R$ and $k \geq 1$. Then $f_4(x) - f_6(x)$ equals

(1) $\frac{1}{4}$

(3) $\frac{1}{6}$

(2) $\frac{1}{12}$

(4) $\frac{1}{3}$

Q68. Let PS be the median of the triangle with vertices $P(2, 2)$, $Q(6, -1)$ and $R(7, 3)$. The equation of the line passing through $(1, -1)$ and parallel to PS is

(1) $4x + 7y + 3 = 0$

(3) $4x - 7y - 11 = 0$

(2) $2x - 9y - 11 = 0$

(4) $2x + 9y + 7 = 0$

Q69. Let a , b , c and d be non-zero numbers. If the point of intersection of the lines

$4ax + 2ay + c = 0$ & $5bx + 2by + d = 0$ lies in the fourth quadrant and is equidistant from the two axes then

(1) $3bc - 2ad = 0$

(3) $2bc - 3ad = 0$

(2) $3bc + 2ad = 0$

(4) $2bc + 3ad = 0$

Q70. Let C be the circle with center at $(1, 1)$ and radius = 1. If T is the circle centered at $(0, y)$, passing through the origin and touching the circle C externally, then the radius of T is equal to

(1) $\frac{1}{2}$

(3) $\frac{\sqrt{3}}{\sqrt{2}}$

(2) $\frac{1}{4}$

(4) $\frac{\sqrt{3}}{2}$

Q71. The locus of the foot of perpendicular drawn from the centre of the ellipse $x^2 + 3y^2 = 6$ on any tangent to it is

(1) $(x^2 + y^2)^2 = 6x^2 + 2y^2$

(3) $(x^2 - y^2)^2 = 6x^2 + 2y^2$

(2) $(x^2 + y^2)^2 = 6x^2 - 2y^2$

(4) $(x^2 - y^2)^2 = 6x^2 - 2y^2$

Q72. $\lim_{x \rightarrow 0} \frac{\sin(\pi \cos^2 x)}{x^2}$ is equal to

(1) $-\pi$

(3) $\frac{\pi}{2}$

(2) π

(4) 1

Q73. The statement $\sim(p \leftrightarrow \sim q)$ is

(1) A tautology

(3) Equivalent to $p \leftrightarrow q$

(2) A fallacy

(4) Equivalent to $\sim p \leftrightarrow q$

Q74. The variance of the first 50 even natural numbers is :

(1) 437

(3) $\frac{833}{4}$

(2) $\frac{437}{4}$

(4) 833

Q75. A bird is sitting on the top of a vertical pole 20 m high and its elevation from a point O on the ground is 45° . It flies off horizontally straight away from the point O . After one second, the elevation of the bird from O is reduced to 30° . Then the speed (in m/s) of the bird is

(1) $20\sqrt{2}$

(3) $40(\sqrt{2} - 1)$

(2) $20(\sqrt{3} - 1)$

(4) $40(\sqrt{3} - \sqrt{2})$

Q76. If $X = \{4^n - 3n - 1 : n \in N\}$ and $Y = \{9(n - 1) : n \in N\}$, where N is the set of natural numbers, then $X \cup Y$ is equal to

(1) X

(3) N

(2) Y

(4) $Y - X$

Q77. If A is a 3×3 non-singular matrix such that $AA' = A'A$ and $B = A^{-1}A'$, then BB' equals, where X' denotes the transpose of the matrix X .

(1) B^{-1}

(3) $I + B$

(2) $(B^{-1})'$

(4) I

Q78.

If $\alpha, \beta \neq 0$, $f(n) = \alpha^n + \beta^n$ and $\begin{vmatrix} 3 & 1+f(1) & 1+f(2) \\ 1+f(1) & 1+f(2) & 1+f(3) \\ 1+f(2) & 1+f(3) & 1+f(4) \end{vmatrix} = K(1-\alpha)^2(1-\beta)^2(\alpha-\beta)^2$, then K is

equal to

(1) 1

(3) $\alpha\beta$

(2) -1

(4) $\frac{1}{\alpha\beta}$

Q79. If g is the inverse of a function f and $f'(x) = \frac{1}{1+x^5}$, then $g'(x)$ is equal to

(1) $\frac{1}{1+\{g(x)\}^5}$

(3) $1+x^5$

(2) $1+\{g(x)\}^5$

(4) $5x^4$

Q80. If f & g are differentiable functions in $[0, 1]$ satisfying $f(0) = 2 = g(1)$, $g(0) = 0$ & $f(1) = 6$, then for some $c \in]0, 1[$

(1) $f'(c) = g'(c)$

(3) $2f'(c) = g'(c)$

(2) $f'(c) = 2g'(c)$

(4) $2f'(c) = 3g'(c)$

Q81. If $x = -1$ and $x = 2$ are extreme points of $f(x) = \alpha \log|x| + \beta x^2 + x$, then

(1) $\alpha = 2, \beta = -\frac{1}{2}$

(3) $\alpha = -6, \beta = \frac{1}{2}$

(2) $\alpha = 2, \beta = \frac{1}{2}$

(4) $\alpha = -6, \beta = -\frac{1}{2}$

Q82. The slope of the line touching both the parabolas $y^2 = 4x$ and $x^2 = -32y$ is

(1) $\frac{1}{8}$

(3) $\frac{1}{2}$

(2) $\frac{2}{3}$

(4) $\frac{3}{2}$

Q83. The integral $\int (1+x-\frac{1}{x})e^{x+\frac{1}{x}}dx$, is equal to

(1) $(x+1)e^{x+\frac{1}{x}} + c$

(3) $(x-1)e^{x+\frac{1}{x}} + c$

(2) $-xe^{x+\frac{1}{x}} + c$

(4) $xe^{x+\frac{1}{x}} + c$

Q84. The integral $\int_0^{\pi} \sqrt{1 + 4 \sin^2 \frac{x}{2} - 4 \sin \frac{x}{2}} dx$ equals

(1) $4\sqrt{3} - 4$

(3) $\pi - 4$

(2) $4\sqrt{3} - 4 - \frac{\pi}{3}$

(4) $\frac{2\pi}{3} - 4 - 4\sqrt{3}$

Q85. The area (in sq. unit) of the region described by $A = \{(x, y) : x^2 + y^2 \leq 1 \text{ and } y^2 \leq 1 - x\}$ is

(1) $\frac{\pi}{2} - \frac{2}{3}$

(3) $\frac{\pi}{2} + \frac{4}{3}$

(2) $\frac{\pi}{2} + \frac{2}{3}$

(4) $\frac{\pi}{2} - \frac{4}{3}$

Q86. Let the population of rabbits surviving at a time t be governed by the differential equation

$\frac{dp(t)}{dt} = \frac{1}{2}\{p(t) - 400\}$. If $p(0) = 100$, then $p(t)$ equals

(1) $600 - 500 e^{\frac{t}{2}}$

(3) $400 - 300 e^{t/2}$

(2) $400 - 300 e^{-\frac{t}{2}}$

(4) $300 - 200 e^{-\frac{t}{2}}$

Q87. If $\left[\vec{a} \times \vec{b} \quad \vec{b} \times \vec{c} \quad \vec{c} \times \vec{a} \right] = \lambda \left[\vec{a} \quad \vec{b} \quad \vec{c} \right]^2$ then λ is equal to

(1) 0

(3) 2

(2) 1

(4) 3

Q88. The image of the line $\frac{x-1}{3} = \frac{y-3}{1} = \frac{z-4}{-5}$ in the plane $2x - y + z + 3 = 0$ is the line

(1) $\frac{x-3}{3} = \frac{y+5}{1} = \frac{z-2}{-5}$

(3) $\frac{x+3}{3} = \frac{y-5}{1} = \frac{z-2}{-5}$

(2) $\frac{x-3}{-3} = \frac{y+5}{-1} = \frac{z-2}{5}$

(4) $\frac{x+3}{-3} = \frac{y-5}{-1} = \frac{z+2}{5}$

Q89. The angle between the lines whose direction cosines satisfy the equations $l + m + n = 0$ and $l^2 = m^2 + n^2$ is

(1) $\frac{\pi}{6}$

(3) $\frac{\pi}{3}$

(2) $\frac{\pi}{2}$

(4) $\frac{\pi}{4}$

Q90. Let A and B be two events such that $P(\overline{A \cup B}) = \frac{1}{6}$, $P(A \cap B) = \frac{1}{4}$ and $P(\overline{A}) = \frac{1}{4}$, where \overline{A} stands for the complement of the event A . Then the events A and B are

(1) Independent but not equally likely.

(3) Mutually exclusive and independent.

(2) Independent and equally likely.

(4) Equally likely but not independent.

ANSWER KEYS

1. (1)	2. (3)	3. (1)	4. (3)	5. (2)	6. (3)	7. (4)	8. (3)
9. (1)	10. (1)	11. (1)	12. (3)	13. (4)	14. (4)	15. (3)	16. (3)
17. (1)	18. (3)	19. (3)	20. (2)	21. (3)	22. (3)	23. (4)	24. (2)
25. (2)	26. (4)	27. (2)	28. (3)	29. (1)	30. (2)	31. (2)	32. (1)
33. (1)	34. (4)	35. (2)	36. (1)	37. (2)	38. (4)	39. (2)	40. (1)
41. (3)	42. (3)	43. (1)	44. (1)	45. (3)	46. (2)	47. (1)	48. (4)
49. (4)	50. (3)	51. (2)	52. (2)	53. (2)	54. (2)	55. (4)	56. (1)
57. (3)	58. (4)	59. (1)	60. (1)	61. (3)	62. (2)	63. (4)	64. (1)
65. (2)	66. (2)	67. (2)	68. (4)	69. (1)	70. (2)	71. (1)	72. (2)
73. (3)	74. (4)	75. (2)	76. (2)	77. (4)	78. (1)	79. (2)	80. (2)
81. (1)	82. (3)	83. (4)	84. (2)	85. (3)	86. (3)	87. (2)	88. (3)
89. (3)	90. (1)						