

Q1. Given, B is magnetic field induction, and μ_0 is the magnetic permeability of vacuum. The dimension of $\frac{B^2}{2\mu_0}$ is:

- (1) MLT^{-2} (2) ML^2T^{-1}
 (3) ML^2T^{-2} (4) $ML^{-1}T^{-2}$

Q2. An elevator in a building can carry a maximum of 10 persons, with the average mass of each person being 68 kg. The mass of the elevator itself is 920 kg and it moves with a constant speed of 3 m/s. The frictional force opposing the motion is 6000 N. If the elevator is moving up with its full capacity, the power delivered by the motor to the elevator ($g = 10\text{m/s}^2$) must be at least:

- (1) 56300W (2) 62360W
 (3) 48000W (4) 66000W

Q3. A mass of 10 kg is suspended by a rope of length 4 m, from the ceiling. A force F is applied horizontally at the mid-point of the rope such that the top half of the rope makes an angle of 45° with the vertical. Then F equals: (Take $g = 10\text{ m s}^{-2}$ and the rope to be massless)

- (1) 100N (2) 90N
 (3) 70N (4) 75N

Q4. Mass per unit area of a circular disc of radius a depends on the distance r from its centre as $\sigma(r) = A + Br$.

The moment of inertia of the disc about the axis, perpendicular to the plane and passing through its centre is:

- (1) $2\pi a^4 \left(\frac{A}{4} + \frac{aB}{5} \right)$ (2) $2\pi a^4 \left(\frac{aA}{4} + \frac{B}{5} \right)$
 (3) $\pi a^4 \left(\frac{A}{4} + \frac{aB}{5} \right)$ (4) $2\pi a^4 \left(\frac{A}{4} + \frac{B}{5} \right)$

Q5. A box weighs 196N on a spring balance at the north pole. Its weight recorded on the same balance if it is shifted to the equator is close to (Take $g = 10\text{ms}^{-2}$ at the north pole and the radius of the earth = 6400km):

- (1) 195.66N (2) 194.32N
 (3) 194.66N (4) 195.32N

Q6. An ideal fluid flows (laminar flow) through a pipe of non-uniform diameter. The maximum and minimum diameters of the pipes are 6.4cm and 4.8cm, respectively. The ratio of the minimum and the maximum velocities of fluid in this pipe is:

- (1) $\frac{9}{16}$ (2) $\frac{\sqrt{3}}{2}$
 (3) $\frac{3}{4}$ (4) $\frac{81}{256}$

Q7. Under an adiabatic process, the volume of an ideal gas gets doubled. Consequently, the mean collision time between the gas molecule changes from τ_1 to τ_2 . If $\frac{C_p}{C_v} = \gamma$ for this gas then a good estimate for $\frac{\tau_2}{\tau_1}$ is given by

- (1) 2 (2) $\frac{1}{2}$
 (3) $\left(\frac{1}{2}\right)^\gamma$ (4) $\left(\frac{1}{2}\right)^{\frac{\gamma+1}{2}}$

Q8. Two ideal Carnot engines operate in cascade (all heat given up by one engine is used by the other engine to produce work) between temperatures, T_1 and T_2 . The temperature of the hot reservoir of the first engine is T_1 and the temperature of the cold reservoir of the second engine is T_2 . T is temperature of the sink of first engine which is also the source for the second engine. How is T related to T_1 and T_2 , if both the engines perform equal amount of work?

- (1) $T = \frac{2T_1T_2}{T_1+T_2}$ (2) $T = \frac{T_1+T_2}{2}$
 (3) $T = \sqrt{T_1T_2}$ (4) $T = 0$

Q9. A stationary observer receives sound from two identical tuning forks, one of which approaches and the other one recedes with the same speed (much less than the speed of sound). The observer hears 2 beats/sec. The oscillation frequency of each tuning fork is $v_0 = 1400$ Hz and the velocity of sound in air is 350 m/s. The speed of each tuning fork is close to:

- (1) $\frac{1}{2}$ m/s (2) 1 m/s
 (3) $\frac{1}{4}$ m/s (4) $\frac{1}{8}$ m/s


Q10. A particle of mass m and charge q has an initial velocity $\vec{v} = v_0\hat{j}$. If an electric field $\vec{E} = E_0\hat{i}$ and magnetic field $\vec{B} = B_0\hat{i}$ act on the particle, its speed will double after a time

- (1) $\frac{2mv_0}{qE_0}$ (2) $\frac{3mv_0}{qE_0}$
 (3) $\frac{\sqrt{3}mv_0}{qE_0}$ (4) $\frac{\sqrt{2}mv_0}{qE_0}$

Q11. In a building there are 15 bulbs of 45W, 15 bulbs of 100W, 15 small fans of 10W and 2 heaters of 1kW. The voltage of electric main supply is 220V. The minimum fuse capacity (rated value) of the building will be:


- (1) 5A (2) 25A
 (3) 15A (4) 20A

Q12. In the figure, potential difference between A and B is:



- (1) 10 V (2) 5 V
 (3) 15 V (4) 0 V

Q13.



The figure gives experimentally measured B vs. H variation in a ferromagnetic material. The retentivity, coercivity and saturation, respectively, of the material are:

(1) 1.5T, 50A/m and 1.0T

(3) 150A/m, 1.0T and 1.5T

(2) 1.5T, 50A/m and 1.0T

(4) 1.0T, 50A/m and 1.5T

Q14. The electric field of a plane electromagnetic wave is given by $\vec{E} = E_0 \frac{\hat{i} + \hat{j}}{\sqrt{2}} \cos(kz + \omega t)$. At $t = 0$, a positively charged particle is at the point $(x, y, z) = (0, 0, \frac{\pi}{k})$. If its instantaneous velocity at $(t = 0)$ is $v_0 \hat{k}$, the force acting on it due to the wave is:

(1) parallel to $\frac{\hat{i} + \hat{j}}{\sqrt{2}}$ (3) antiparallel to $\frac{\hat{i} + \hat{j}}{\sqrt{2}}$

(2) zero

(4) parallel to \hat{k}

Q15. A planar loop of wire rotates in a uniform magnetic field. Initially, at $t = 0$, the plane of the loop is perpendicular to the magnetic field. If it rotates with a period of 10s about an axis in its plane then the magnitude of induced emf will be maximum and minimum, respectively at:

(1) 2.5 s and 7.5 s

(3) 5.0 s and 7.5 s

(2) 2.5 s and 5.0 s

(4) 5.0 s and 10.0 s

Q16. An emf of 20 V is applied at time $t = 0$ to a circuit containing in series 10 mH inductor and 5Ω resistor. The ratio of the currents at time $t = \infty$ and at $t = 40$ s is close to: (Take $e^2 = 7.389$)

(1) 1.06

(3) 1.46

(2) 1.15

(4) 0.84

Q17. A thin lens made of glass (refractive index = 1.5) of focal length $f = 16$ cm is immersed in a liquid of refractive index 1.42. If its focal length in liquid is f_l , then the ratio f_l/f is closest to the integer:

(1) 1

(3) 5

(2) 9

(4) 17

Q18. In a Young's double slit experiment, the separation between the slits is 0.15mm. In the experiment, a source of light of wavelength 589nm is used and the interference pattern is observed on a screen kept 1.5m away. The separation between the successive bright fringes on the screen is:

(1) 6.9mm

(3) 5.9mm

(2) 3.9mm

(4) 4.9mm

Q19. An electron (of mass m) and a photon have the same energy E in the range of a few eV. The ratio of the de-Broglie wavelength associated with the electron and the wavelength of the photon is (c = speed of light in vacuum)

(1) $\frac{1}{c} \left(\frac{2E}{m} \right)^{1/2}$ (3) $\frac{1}{c} \left(\frac{E}{2m} \right)^{1/2}$ (2) $c(2mE)^{1/2}$ (4) $\left(\frac{E}{2m} \right)^{1/2}$

Q20. The activity of a radioactive sample falls from 700s^{-1} to 500s^{-1} in 30 minutes. Its half life is close to:

(1) 72min

(3) 66min


(2) 62min

(4) 52min

Q21. The sum of two forces \vec{P} and \vec{Q} is \vec{R} such that $|\vec{R}| = |\vec{P}|$. Find the angle between resultant of $2\vec{P}$ and \vec{Q} and \vec{Q}

, _____

Q22.



Consider a uniform cubical box of side a on a rough floor that is to be moved by applying minimum possible force F at a point b above its centre of mass (see figure). If the coefficient of friction is $\mu = 0.4$, the maximum possible value of $100 \times \frac{b}{a}$ for a box not to topple before moving is _____

Q23. M grams of steam at 100°C is mixed with 200g of ice at its melting point in a thermally insulated container. If it produces liquid water at 40°C [heat of vaporization of water is 540cal/g and heat of fusion of ice is 80cal/g], the value of M is _____

Q24. A 60pF capacitor is fully charged by a 20V supply. It is then disconnected from the supply and is connected to another uncharged 60pF capacitor in parallel. The electrostatic energy that is lost in this process by the time the charge is redistributed between them is (in nJ) _____

Q25. The balancing length for a cell is 560cm in a potentiometer experiment. When an external resistance of 10Ω is connected in parallel to the cell, the balancing length changes by 60cm . If the internal resistance of the cell is $\frac{n}{10}\Omega$, where n is an integer then value of n is _____

Q26. The ammonia (NH_3) released on quantitative reaction of 0.6g urea (NH_2CONH_2) with sodium hydroxide (NaOH) can be neutralized by

- | | |
|--|--|
| (1) 200 ml of 0.4 NHCl | (2) 200 ml of 0.2 NHCl |
| (3) 100 ml of 0.2 NHCl | (4) 100 ml of 0.1 NHCl |


Q27. Within each pair of elements F and Cl , S and Se , and Li and Na , respectively, the elements that release more energy upon an electron gain are

- | | |
|---|--|
| (1) Cl , Se and Na | (2) Cl , S and Li |
| (3) F , S and Li | (4) F , Se and Na |

Q28. The bond order and the magnetic characteristic of CN^- are

- | | |
|----------------------------------|-----------------------------------|
| (1) $2\frac{1}{2}$, diamagnetic | (2) 3 , diamagnetic |
| (3) 3 , paramagnetic | (4) $2\frac{1}{2}$, paramagnetic |

Q29. Identify the correct labels of A , B and C in the following graph from the options given below:



Root mean square speed (V_{rms}); most probable speed (V_{mp}); Average speed (V_{av})

(1) $A - V_{mp}; B - V_{rms}; C - V_{av}$

(3) $A - V_{rms}; B - V_{mp}; C - V_{av}$

(2) $A - V_{av}; B - V_{rms}; C - V_{mp}$

(4) $A - V_{mp}; B - V_{av}; C - V_{rms}$

Q30. The redox reaction among the following is

(1) formation of ozone from atmospheric oxygen in the presence of sunlight

(3) reaction of H_2SO_4 with NaOH

(2) reaction of $[Co(H_2O)_6]Cl_3$ with $AgNO_3$

(4) combination of dinitrogen with dioxygen at 2000K

Q31. Among statements (a) – (d), the correct ones are:

(a) Decomposition of hydrogen peroxide gives di-oxygen.

(b) Like hydrogen peroxide, compounds, such as $KClO_3$, $Pb(NO_3)_2$ and $NaNO_3$ When heated liberate di-oxygen.

(c) 2-Ethylanthraquinone is useful for the industrial preparation of hydrogen peroxide.

(d) Hydrogen peroxide is used for the manufacture of sodium perborate.


(1) (a), (b), (c) and (d)

(3) (a), (c), and (d) only


(2) (a), (b), and (c) only

(4) (a) and (c) only


Q32. The correct order of stability for the following alkoxides is:



(A)



(B)



(C)

(1) $(B) > (A) > (C)$

(3) $(C) > (A) > (B)$

(2) $(C) > (B) > (A)$

(4) $(B) > (C) > (A)$

Q33. A chromatography column, packed with silica gel as stationary phase, was used to separate a mixture of compounds consisting of (A) benzanilide (B) aniline and (C) acetophenone. When the column is eluted with a mixture of solvents, hexane:ethyl acetate (20 : 80), the sequence of obtained compounds is:

(1) (B), (C) and (A)

(3) (C), (A) and (B)

(2) (B), (A) and (C)

(4) (A), (B) and (C)

Q34. Two open beakers one containing a solvent and the other containing a mixture of that solvent with a non volatile solute are together sealed in a container. Over time:

(1) the volume of the solution increases and the volume of the solvent decreases

(3) the volume of the solution and the solvent does not change the volume of the solution does not

(2) the volume of the solution decreases and the volume of the solvent increases

(4) change and the volume of the solvent decreases

Q35. The equation that is incorrect is:

- (1) $(\Delta_m^0)_{\text{KBr}} - (\Delta_m^0)_{\text{NaCl}} = (\Delta_m^0)_{\text{KBr}} - (\Delta_m^0)_{\text{KCl}}$
 (2) $(\Delta_m^0)_{\text{KG}} - (\Delta_m^0)_{\text{KQ}} = (\Delta_m^0)_{\text{NaG}} - (\Delta_m^0)_{\text{NaQ}}$
 (3) $(\Delta_m^0)_{\text{H}_2\text{O}} = (\Delta_m^0)_{\text{HCl}} + (\Delta_m^0)_{\text{NaOH}} - (\Delta_m^0)_{\text{NaCl}}$
 (4) $(\Delta_m^0)_{\text{NaBr}} - (\Delta_m^0)_{\text{NaI}} = (\Delta_m^0)_{\text{KBr}} - (\Delta_m^0)_{\text{KI}}$

Q36. For the reaction $2\text{H}_2(\text{g}) + 2\text{NO}(\text{g}) \rightarrow \text{N}_2(\text{g}) + 2\text{H}_2\text{O}(\text{g})$ the observed rate expression is, $\text{rate} = k_f[\text{NO}]^2[\text{H}_2]$.

The rate expression for the reverse reaction is:

- (1) $k_b[\text{N}_2][\text{H}_2\text{O}]^2$ (2) $k_b[\text{N}_2][\text{H}_2\text{O}]^2/[\text{NO}]$
 (3) $k_b[\text{N}_2][\text{H}_2\text{O}]$ (4) $k_b[\text{N}_2][\text{H}_2\text{O}]^2/[\text{H}_2]$

Q37. The refining method used when the metal and the impurities have low and high melting temperatures, respectively, is

- (1) liquation. (2) vapour phase refining.
 (3) zone refining. (4) distillation.

Q38. In the following reactions, products (A) and (B), respectively, are:

$\text{NaOH} + \text{Cl}_2 \rightarrow (\text{A}) + \text{side products}$
 (hot and conc.)

$\text{Ca}(\text{OH})_2 + \text{Cl}_2 \rightarrow (\text{B}) + \text{side products}$
 (dry)

- (1) NaClO_3 and $\text{Ca}(\text{OCl})_2$ (2) NaClO_3 and $\text{Ca}(\text{ClO}_3)_2$
 (3) NaOCl and $\text{Ca}(\text{OCl})_2$ (4) NaOCl and $\text{Ca}(\text{ClO}_3)_2$

Q39. The number of possible optical isomers for the complexes MA_2B_2 with sp^3 and dsp^2 hybridized metal atom, respectively, is:

Note: A and B are unidentate neutral and unidentate monoanionic ligands, respectively.


- (1) 0 and 2 (2) 2 and 2
 (3) 0 and 0 (4) 0 and 1

Q40. Among the statements (a) – (d), the incorrect ones are:


- (a) Octahedral $\text{Co}(\text{III})$ complexes with strong field ligands have very high magnetic moments
 (b) When $\Delta_0 < P$, the d-electron configuration of $\text{Co}(\text{III})$ in an octahedral complex is $t_{eg}^4 e_g^2$
 (c) Wavelength of light absorbed by $[\text{Co}(\text{en})_3]^{3+}$ is lower than that of $[\text{CoF}_6]^{3-}$
 (d) If the Δ_0 for an octahedral complex of $\text{Co}(\text{III})$ is $18,000\text{cm}^{-1}$, the Δ_t for its tetrahedral complex with the same ligand will be $16,000\text{cm}^{-1}$.

- (1) (a) and (d) only (2) (c) and (d) only
 (3) (a) and (b) only (4) (b) and (c) only

Q41. For the following reactions



where,



k_s and k_e , are respectively, the rate constants for substitution and elimination, and $\mu = \frac{k_s}{k_e}$, the correct option is _____









(1) $\mu_B > \mu_A$ and $k_e(\text{A}) > k_e(\text{B})$

(2) $\mu_A > \mu_B$ and $k_e(\text{B}) > k_e(\text{A})$

(3) $\mu_B > \mu_A$ and $k_e(\text{B}) > k_e(\text{A})$

(4) $\mu_A > \mu_B$ and $k_e(\text{A}) > k_e(\text{B})$

Q42. Consider the following reactions:



Which of these reactions are possible?






(1) (a) and (b)

(3) (b), (c) and (d)


(2) (a) and (d)

(4) (b) and (d)




Q43. In the following reaction sequence, structures of A and B are, respectively



Q44. In the following reaction sequence,



the major product B is



Q45. Which of the following statements is correct?

(1) Gluconic acid can form cyclic (acetal/hemiacetal) structure

(3) Gluconic acid is a partial oxidation product of glucose

(2) Gluconic acid is a dicarboxylic acid

(4) Gluconic acid is obtained by oxidation of glucose with HNO_3

- Q46.** The standard heat of formation ($\Delta_f H_{298}^0$) of ethane (in kJ/mol), if the heat of combustion of ethane, hydrogen and graphite are -1560 , -393.5 and -286 kJ/mol, respectively is _____
- Q47.** 3 g of acetic acid is added to 250 mL of 0.1 M HCl and the solution made up to 500 mL. To 20 mL of this solution $\frac{1}{2}$ mL of 5M NaOH is added. The pH of the solution is _____
[Given: pK_a of acetic acid = 4.75, molar mass of acetic acid 60 g/mol, $\log 3 = 0.4771$, Neglect any changes in volume]
- Q48.** Consider the following reactions:

$$\text{NaCl} + \text{K}_2\text{Cr}_2\text{O}_7 + \text{H}_2\text{SO}_4 \rightarrow (\text{A}) + \text{side products}$$
(Conc.)

$$(\text{A}) + \text{NaOH} \rightarrow (\text{B}) + \text{side products}$$

$$(\text{B}) + \text{H}_2\text{SO}_4 + \text{H}_2\text{O}_2 \rightarrow (\text{C}) + \text{side products}$$
(dilute)
 The sum of the total number of atoms in one molecule each of (A), (B) and (C) is _____
- Q49.** The flocculation value of HCl for arsenic sulphide sol is 30 mmol L^{-1} . If H_2SO_4 is used for the flocculation of arsenic sulphide, the amount, in grams, of H_2SO_4 in 250ml required for the above purpose is _____
(molecular mass of $\text{H}_2\text{SO}_4 = 98 \text{ g/mol}$)
- Q50.** The number of sp^2 hybridised carbons present in "Aspartame" is _____
- Q51.** Let α and β be the roots of the equation $x^2 - x - 1 = 0$. If $p_k = (\alpha)^k + (\beta)^k$, $k \geq 1$, then which one of the following statements is not true?
 (1) $p_3 = p_5 - p_4$ (2) $p_5 = 11$
 (3) $(p_1 + p_2 + p_3 + p_4 + p_5) = 26$ (4) $p_5 = p_2 \cdot p_3$
- Q52.** If $\frac{3+i\sin\theta}{4-i\cos\theta}$, $\theta \in [0, 2\pi]$, is a real number, then an argument of $\sin\theta + i\cos\theta$ is
 (1) $\pi - \tan^{-1}\left(\frac{4}{3}\right)$ (2) $\pi - \tan^{-1}\left(\frac{3}{4}\right)$
 (3) $-\tan^{-1}\left(\frac{3}{4}\right)$ (4) $\tan^{-1}\left(\frac{4}{3}\right)$
- Q53.** Let a_1, a_2, a_3, \dots , be a G. P. such that $a_1 < 0$, $a_1 + a_2 = 4$ and $a_3 + a_4 = 16$. If $\sum_{i=1}^9 a_i = 4\lambda$, then λ , is equal to.
 (1) -513 (2) -171
 (3) 171 (4) $\frac{511}{3}$
- Q54.** If the sum of the first 40 terms of the series, $3 + 4 + 8 + 9 + 13 + 14 + 18 + 19 + \dots$ is $(102)m$, then m is equal to
 (1) 20 (2) 25
 (3) 5 (4) 10
- Q55.** The coefficient of x^7 in the expression $(1+x)^{10} + x(1+x)^9 + x^2(1+x)^8 + \dots + x^{10}$, is
 (1) 210 (2) 330
 (3) 120 (4) 420
- Q56.** The number of ordered pairs (r, k) for which $6 \cdot {}^{35}C_r = (k^2 - 3) \cdot {}^{36}C_{r+1}$, where k is an integer is _____

(1) 3

(3) 6

(2) 2

(4) 4

Q57. The locus of the mid-points of the perpendiculars drawn from points on the line $x = 2y$, to the line $x = y$, is.

(1) $2x - 3y = 0$ (3) $3x - 2y = 0$ (2) $5x - 7y = 0$ (4) $7x - 5y = 0$

Q58. Let the tangents drawn from the origin to the circle, $x^2 + y^2 - 8x - 4y + 16 = 0$ touch it at the points A and B . Then $(AB)^2$ is equal to

(1) $\frac{52}{5}$ (3) $\frac{64}{5}$ (2) $\frac{56}{5}$ (4) $\frac{32}{5}$

Q59. If $3x + 4y = 12\sqrt{2}$ is a tangent to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{9} = 1$ for some $a \in R$, then the distance between the foci of the ellipse is

(1) $2\sqrt{7}$ (3) $2\sqrt{5}$

(2) 4

(4) $2\sqrt{2}$

Q60. Let A, B, C and D be four non-empty sets. The contrapositive statement of "If $A \subseteq B$ and $B \subseteq D$, then $A \subseteq C$ " is

(1) If $A \not\subseteq C$, then $A \subseteq B$ and $B \subseteq D$ (3) If $A \not\subseteq C$, then $A \not\subseteq B$ and $B \subseteq D$ (2) If $A \subseteq C$, then $B \subset A$ and $D \subset B$ (4) If $A \not\subseteq C$, then $A \not\subseteq B$ or $B \not\subseteq D$

Q61. Let $A = [a_{ij}]$ and $B = [b_{ij}]$ be two 3×3 real matrices such that $b_{ij} = (3)^{(i+j-2)} a_{ij}$, where $i, j = 1, 2, 3$. If the determinant of B is 81, then determinant of A is

(1) $\frac{1}{3}$ (3) $\frac{1}{81}$

(2) 3

(4) $\frac{1}{9}$

Q62. Let $y = y(x)$ be a function of x satisfying $y\sqrt{1-x^2} = k - x\sqrt{1-y^2}$ where k is a constant and $y(\frac{1}{2}) = -\frac{1}{4}$. Then $\frac{dy}{dx}$ at $x = \frac{1}{2}$, is equal to

(1) $-\frac{\sqrt{5}}{4}$ (3) $\frac{2}{\sqrt{5}}$ (2) $-\frac{\sqrt{5}}{2}$ (4) $\frac{\sqrt{5}-2}{2}$

Q63. The value of c , in the Lagrange's mean value theorem for the function $f(x) = x^3 - 4x^2 + 8x + 11$, when $x \in [0, 1]$, is

(1) $\frac{4-\sqrt{5}}{3}$ (3) $\frac{2}{3}$ (2) $\frac{4-\sqrt{7}}{3}$ (4) $\frac{\sqrt{7}-2}{3}$

Q64. Let $f(x)$ be a polynomial of degree 5 such that $x = \pm 1$ are its critical points. If $\lim_{x \rightarrow 0} \left(2 + \frac{f(x)}{x^3} \right) = 4$, then which one of the following is not true?

(1) f is an odd function

(3) $x = 1$ is a point of local minimum and $x = -1$ is a point of local maximum

(2) $f(1) - 4f(-1) = 4$. $x = 1$ is a point of maximum and $x = -1$

(4) $x = 1$ is a point of local maxima of f

Q65. The value of α for which $4\alpha \int_{-1}^2 e^{-\alpha|x|} dx = 5$, is

- (1) $\log_e 2$ (2) $\log_e \left(\frac{3}{2}\right)$
 (3) $\log_e \sqrt{2}$ (4) $\log_e \left(\frac{4}{3}\right)$

Q66. If θ_1 and θ_2 be respectively the smallest and the largest values of θ in $(0, 2\pi) - \{\pi\}$ which satisfy the equation,

$$2\cot^2\theta - \frac{5}{\sin\theta} + 4 = 0, \text{ then } \int_{\theta_1}^{\theta_2} \cos^2 3\theta d\theta \text{ is equal to:}$$

- (1) $\frac{\pi}{3}$ (2) $\frac{2\pi}{3}$
 (3) $\frac{\pi}{3} + \frac{1}{6}$ (4) $\frac{\pi}{9}$

Q67. The area (in sq. units) of the region $\{(x, y) \in R^2 | 4x^2 \leq y \leq 8x + 12\}$ is

- (1) $\frac{125}{3}$ (2) $\frac{128}{3}$
 (3) $\frac{124}{3}$ (4) $\frac{127}{3}$

Q68. Let $y = y(x)$ be the solution curve of the differential equation, $(y^2 - x) \frac{dy}{dx} = 1$, satisfying $y(0) = 1$. This curve intersects the X -axis at a point whose abscissa is

- (1) $2 - e$ (2) $-e$
 (3) 2 (4) $2 + e$

Q69. Let \vec{a}, \vec{b} and \vec{c} , be three unit vectors such that $\vec{a} + \vec{b} + \vec{c} = \vec{0}$. If $\lambda = \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}$ and $\vec{d} = \vec{a} \times \vec{b} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a}$, then the order pair, (λ, \vec{d}) , is equal to.

- (1) $\left(\frac{3}{2}, 3\vec{a} \times \vec{c}\right)$ (2) $\left(-\frac{3}{2}, 3\vec{c} \times \vec{b}\right)$
 (3) $\left(\frac{3}{2}, 3\vec{b} \times \vec{c}\right)$ (4) $\left(-\frac{3}{2}, 3\vec{a} \times \vec{b}\right)$

Q70. In a workshop, there are five machines and the probability of any one of them to be out of service on a day is $\frac{1}{4}$. If the probability that at most two machines will be out of service on the same day is $\left(\frac{3}{4}\right)^3 k$, then k is equal to

- (1) $\frac{17}{8}$ (2) $\frac{17}{4}$
 (3) $\frac{17}{2}$ (4) 4

Q71. If the mean and variance of eight numbers 3, 7, 9, 12, 13, 20, x and y be 10 and 25 respectively, then $x \cdot y$ is equal to

Q72. Let $X = \{n \in N : 1 \leq n \leq 50\}$. If $A = \{n \in X : n \text{ is a multiple of } 2\}$ and

$B = \{n \in X : n \text{ is a multiple of } 7\}$, then the number of elements in the smallest subset of X , containing both A and B , is.

Q73. If the system of linear equations,

$$x + y + z = 6$$

$$x + 2y + 3z = 10$$

$$3x + 2y + \lambda z = \mu$$

has more than two solutions, then $\mu - \lambda^2$, is equal to.

Q74. If the function f defined on $(-\frac{1}{3}, 1/3)$ by $f(x) = \begin{cases} \frac{1}{x} \log_e \left(\frac{1+3x}{1-2x} \right), & \text{when } x \neq 0 \\ k, & \text{when } x = 0 \end{cases}$, is continuous, then k is equal to.

Q75. If the foot of the perpendicular drawn from the point $(1, 0, 3)$ on a line passing through $(\alpha, 7, 1)$ is $(\frac{5}{3}, \frac{7}{3}, \frac{17}{3})$, then α is equal to

ANSWER KEYS

1. (4)	2. (4)	3. (1)	4. (1)	5. (4)	6. (1)	7. (4)	8. (2)
9. (3)	10. (3)	11. (4)	12. (1)	13. (4)	14. (3)	15. (2)	16. (1)
17. (2)	18. (3)	19. (3)	20. (2)	21. (90)	22. (50)	23. (40)	24. (6)
25. (12)	26. (3)	27. (2)	28. (2)	29. (4)	30. (4)	31. (1)	32. (2)
33. (3)	34. (1)	35. (1)	36. (4)	37. (1)	38. (1)	39. (3)	40. (1)
41. (2)	42. (4)	43. (3)	44. (1)	45. (3)	46. (-192)	47. (5.22)	48. (18)
49. (0.36)	50. (9)	51. (4)	52. (1)	53. (2)	54. (1)	55. (2)	56. (4)
57. (2)	58. (3)	59. (1)	60. (4)	61. (4)	62. (2)	63. (2)	64. (4)
65. (1)	66. (1)	67. (2)	68. (1)	69. (4)	70. (1)	71. (54)	72. (29)
73. (13)	74. (5)	75. (4)					