

# JEE-MAIN EXAMINATION – JANUARY 2025

(HELD ON TUESDAY 28<sup>th</sup> JANUARY 2025)

TIME : 9:00 AM TO 12:00 NOON

## MATHEMATICS

### SECTION-A

1. The number of different 5 digit numbers greater than 50000 that can be formed using the digits 0, 1, 2, 3, 4, 5, 6, 7, such that the sum of their first and last digits should not be more than 8, is

- (1) 4608                      (2) 5720  
(3) 5719                      (4) 4607

**Ans. (4)**

**Sol.** Case I    5 \_ \_ \_ 0

Case II    5 \_ \_ \_ 1

5        2

5        3

6        0

6        1

6        2

7        0

Case IX    7 \_ \_ \_ 1

$9 \times (8 \times 8 \times 8) = 4608$  but 50000 is not included, so total numbers  $4608 - 1 = 4607$


2. Let ABCD be a trapezium whose vertices lie on the parabola  $y^2 = 4x$ . Let the sides AD and BC of the trapezium be parallel to y-axis. If the diagonal

AC is of length  $\frac{25}{4}$  and it passes through the point

(1, 0), then the area of ABCD is :

- (1)  $\frac{75}{4}$                       (2)  $\frac{25}{2}$   
(3)  $\frac{125}{8}$                       (4)  $\frac{75}{8}$

**Ans. (1)**



$$A(at_1^2, 2at_1) \text{ \& \; } C\left(\frac{a}{t_1^2}, -\frac{2a}{t_1}\right)$$

$$\text{Length AC} = a \left( t_1 + \frac{1}{t_1} \right)^2 = \frac{25}{4}, \quad t_1 + \frac{1}{t_1} = \pm \frac{5}{2}$$

$$\Rightarrow t_1 = 2 \text{ or } \frac{1}{2}, \quad A\left(\frac{1}{2}, 1\right), D\left(\frac{1}{4}, -1\right), B(4, 4), C(4, -4)$$

$$\text{So, area of trapezium} = \frac{1}{2}(8+2)\left(4 - \frac{1}{4}\right) = \frac{75}{4}$$

3. Two number  $k_1$  and  $k_2$  are randomly chosen from the set of natural numbers. Then, the probability that the value of  $i^{k_1} + i^{k_2}$ , ( $i = \sqrt{-1}$ ) is non-zero, equals

- (1)  $\frac{1}{2}$                       (2)  $\frac{1}{4}$   
(3)  $\frac{3}{4}$                       (4)  $\frac{2}{3}$

**Ans. (3)**

**Sol.**  $i^{k_1} + i^{k_2} \neq 0 \quad i^{k_1} \rightarrow 4 \text{ option for } i, -1, -i, 1$

Total cases  $\Rightarrow 4 \times 4 = 16$

Unfavourable cases  $\Rightarrow i^{k_1} + i^{k_2} = 0$

$$\begin{Bmatrix} 1, -1 \\ -1, 1 \\ i, -i \\ -i, i \end{Bmatrix}$$

$$4 \text{ Cases} \Rightarrow \text{Probability} = \frac{16-4}{16} = \frac{3}{4}$$

4. If  $f(x) = \frac{2^x}{2^x + \sqrt{2}}$ ,  $x \in \mathbb{R}$ , then  $\sum_{k=1}^{81} f\left(\frac{k}{82}\right)$  is equal

to :

- (1) 41                      (2)  $\frac{81}{2}$   
(3) 82                      (4)  $81\sqrt{2}$

**Ans. (2)**



**Sol.**  $f(x) = \frac{2^x}{2^x + \sqrt{2}}$

$$f(x) + f(1-x) = \frac{2^x}{2^x + \sqrt{2}} + \frac{2^{1-x}}{2^{1-x} + \sqrt{2}}$$

$$= \frac{2^x}{2^x + \sqrt{2}} + \frac{2}{2 + \sqrt{2} 2^x} = \frac{2^x + \sqrt{2}}{2^x + \sqrt{2}} = 1$$

Now,  $\sum_{k=1}^{81} f\left(\frac{k}{82}\right) = f\left(\frac{1}{82}\right) + f\left(\frac{2}{82}\right) + \dots + f\left(\frac{81}{82}\right)$

$$= f\left(\frac{1}{82}\right) + f\left(\frac{2}{82}\right) + \dots + f\left(1 - \frac{2}{82}\right) + f\left(1 - \frac{1}{82}\right)$$

$$= \left[ f\left(\frac{1}{82}\right) + f\left(1 - \frac{1}{82}\right) \right] + \left[ f\left(\frac{2}{82}\right) + f\left(1 - \frac{2}{82}\right) \right] + \dots + 40 \text{ cases} + f\left(\frac{41}{82}\right)$$

$$= (1+1+\dots 40 \text{ times}) + \frac{2^{1/2}}{2^{1/2} + 2^{1/2}}$$

$$40 + \frac{1}{2} = \frac{81}{2}$$

5. Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a function defined by

$$f(x) = (2+3a)x^2 + \left(\frac{a+2}{a-1}\right)x + b, a \neq 1. \text{ If}$$

$$f(x+y) = f(x) + f(y) + 1 - \frac{2}{7}xy, \text{ then the value of}$$

$$28 \sum_{i=1}^5 |f(i)| \text{ is:}$$

(1) 715

(2) 735

(3) 545

(4) 675

**Ans. (4)**

**Sol.**  $f(x) = (3a+2)x^2 + \left(\frac{a+2}{a-1}\right)x + b$

$$f\left(x + \frac{1}{2}\right) = f(x) + f(y) + 1 - \frac{2}{7}xy \dots (1)$$

In (1) Put  $x = y = 0 \Rightarrow f(0) = 2f(0) + 1 \Rightarrow f(0) = -1$

So,  $f(0) = 0 + 0 + b = -1 \Rightarrow b = -1$

In (1) Put  $y = -x \Rightarrow f(0) = f(x) + f(-x) + 1 + \frac{2}{7}x^2$

$$-1 = 2(3a+2)x^2 + 2b + 1 + \frac{2}{7}x^2$$

$$-1 = \left(2(3a+2) + \frac{2}{7}\right)x^2 + 1 - 2$$

$$\Rightarrow 6a + 4 + \frac{2}{7} = 0$$

$$a = -\frac{5}{7}$$

So  $f(x) = -\frac{1}{7}x^2 - \frac{3}{4}x - 1$

$$\Rightarrow |f(x)| = \frac{1}{28} |4x^2 + 21x + 28|$$

Now,  $28 \sum_{i=1}^5 |f(i)| = 28(|f(1)| + |f(2)| + \dots + |f(5)|)$

$$28 \cdot \frac{1}{28} \cdot 675 = 675$$

6. Let  $A(x, y, z)$  be a point in  $xy$ -plane, which is equidistant from three points  $(0, 3, 2)$ ,  $(2, 0, 3)$  and  $(0, 0, 1)$ .

Let  $B = (1, 4, -1)$  and  $C = (2, 0, -2)$ . Then among the statements

(S1) :  $\Delta ABC$  is an isosceles right angled triangle and

(S2) : the area of  $\Delta ABC$  is  $\frac{9\sqrt{2}}{2}$ .

(1) both are true

(2) only (S1) is true

(3) only (S2) is true

(4) both are false

**Ans. (2)**

**Sol.**  $A(x, y, z)$  Let  $P(0, 3, 2)$ ,  $Q(2, 0, 3)$ ,  $R(0, 0, 1)$

$$AP = AQ = AR$$

$$x^2 + (y-3)^2 + (z-2)^2 = (x-2)^2 + y^2 + (z-3)^2 = x^2 + y^2 + (z-1)^2$$

In  $xy$  plane  $z = 0$

$$\text{So, } x^2 - 4x + 4 + y^2 + 9 = x^2 + y^2 + 1$$

$$\Rightarrow y = 2$$

$$x = 3$$

$$9 + y^2 - 6y + 9 + 4 = x^2 + y^2 + 1$$

So,  $A(3, 2, 0)$  also  $B(1, 4, -1)$  &  $C(2, 0, -2)$

$$\text{Now } AB = \sqrt{4+4+1} = 3$$



$$AC = \sqrt{1+4+4} = 3$$

$$BC = \sqrt{1+16+1} = \sqrt{18}$$

$$AB = AC$$

$$\text{isosceles } \Delta \text{ \& } AB^2 + AC^2 = BC^2$$

right angle  $\Delta$

$$\text{Area of } \Delta ABC = \frac{1}{2} \times \text{base} \times \text{height}$$

$$\frac{1}{2} \times 3 \times 3 = \frac{9}{2}$$

So only  $S_1$  is true

7. The relation  $R = \{(x, y) : x, y \in Z \text{ and } x + y \text{ is even}\}$  is :

- (1) reflexive and transitive but not symmetric
- (2) reflexive and symmetric but not transitive
- (3) an equivalence relation
- (4) symmetric and transitive but not reflexive

**Ans. (3)**

**Sol.**  $R = \{(x, y) : x, y \in Z \text{ and } x + y \text{ is even}\}$

reflexive  $x + x = 2x$  even

symmetric of  $x + y$  is even, then  $(y + x)$  is also even

transitive of  $x + y$  is even &  $y + z$  is even then  $x + z$  is also even


So, relation is an equivalence relation.

8. Let the equation of the circle, which touches x-axis at the point  $(a, 0)$ ,  $a > 0$  and cuts off an intercept of length  $b$  on y-axis be  $x^2 + y^2 - \alpha x + \beta y + \gamma = 0$ . If the circle lies below x-axis, then the ordered pair  $(2a, b^2)$  is equal to :

- (1)  $(\alpha, \beta^2 + 4\gamma)$
- (2)  $(\gamma, \beta^2 - 4\alpha)$
- (3)  $(\gamma, \beta^2 + 4\alpha)$
- (4)  $(\alpha, \beta^2 - 4\gamma)$

**Ans. (4)**

**Sol.**



$$\text{By pythagorus } r^2 = a^2 + \frac{b^2}{4} = p^2$$

$$r = \sqrt{\frac{4a^2 + b^2}{4}}$$

$$\text{Equation of circle is } (x - \alpha)^2 + (y - \beta)^2 = r^2$$

$$x^2 + y^2 - 2\alpha x - 2\beta y + \alpha^2 + \beta^2 - r^2 = 0$$

$$\text{comparision } x^2 + y^2 - \alpha x + \beta y + r = 0$$

$$-\alpha = -2a, \beta = -2p, r = a^2$$

$$\Rightarrow 2a = \alpha, 4a^2 + b^2 = 4p^2$$

$$\alpha^2 + b^2 = 4p^2$$

$$\alpha^2 + b^2 = \beta^2$$

$$\text{So, } (2a, b^2) = (\alpha, \beta^2 - 4r)$$

9. Let  $\langle a_n \rangle$  be a sequence such that  $a_0 = 0$ ,  $a_1 = \frac{1}{2}$  and

$$2a_{n+2} = 5a_{n+1} - 3a_n, n = 0, 1, 2, 3, \dots. \text{ Then } \sum_{k=1}^{100} a_k$$

is equal to :

$$(1) 3a_{99} - 100 \quad (2) 3a_{100} - 100$$

$$(3) 3a_{100} + 100 \quad (4) 3a_{99} + 100$$

**Ans. (2)**

$$\text{Sol. } a_0 = 0, a_1 = \frac{1}{2}$$

$$2a_{n+2} = 5a_{n+1} - 3a_n$$

$$2x^2 - 5x + 3 = 0 \Rightarrow x = 1, 3/2$$

$$\therefore a_n = A(1)^n + B\left(\frac{3}{2}\right)^n$$

$$n = 0 \quad 0 = A + B \quad \left. \begin{array}{l} \\ \end{array} \right\} A = -1$$

$$n = 1 \quad \frac{1}{2} = A + \frac{3}{2}B \quad \left. \begin{array}{l} \\ \end{array} \right\} B = 1$$

$$\Rightarrow a_n = -1 + \left(\frac{3}{2}\right)^n$$

$$\sum_{k=1}^{100} a_k = \sum_{k=1}^{100} (-1) + \left(\frac{3}{2}\right)^k$$



$$= -100 + \frac{\left(\frac{3}{2}\right)\left(\left(\frac{3}{2}\right)^{100} - 1\right)}{\frac{3}{2} - 1}$$

$$= -100 + 3\left(\left(\frac{3}{2}\right)^{100} - 1\right)$$

$$= 3(a_{100}) - 100$$

10.  $\cos\left(\sin^{-1}\frac{3}{5} + \sin^{-1}\frac{5}{13} + \sin^{-1}\frac{33}{65}\right)$  is equal to :

(1) 1

(2) 0

(3)  $\frac{33}{65}$

(4)  $\frac{32}{65}$

Ans. (2)

Sol.  $\cos\left(\sin^{-1}\frac{3}{5} + \sin^{-1}\frac{5}{13} + \sin^{-1}\frac{33}{65}\right)$

$$\cos\left(\tan^{-1}\frac{3}{4} + \tan^{-1}\frac{5}{12} + \tan^{-1}\frac{33}{56}\right)$$

$$\cos\left(\tan^{-1}\left(\frac{\frac{3}{4} + \frac{5}{12}}{1 + \frac{3}{4} \cdot \frac{5}{12}}\right) + \tan^{-1}\frac{33}{56}\right)$$

$$\cos\left(\tan^{-1}\frac{56}{33} + \cot^{-1}\frac{56}{33}\right)$$

$$\cos\left(\frac{\pi}{2}\right) = 0$$

11. Let  $T_r$  be the  $r^{\text{th}}$  term of an A.P. If for some  $m$ ,

$$T_m = \frac{1}{25}, T_{25} = \frac{1}{20} \text{ and } 20 \sum_{r=1}^{25} T_r = 13, \text{ then}$$

$$5m \sum_{r=m}^{2m} T_r \text{ is equal to :}$$

(1) 112

(2) 126

(3) 98

(4) 142

Ans. (2)

Sol.  $T_m = \frac{1}{25}, T_{25} = \frac{1}{20}, 20 \sum_{r=1}^{25} T_r = 13$

$$T_m = a + (m-1)d = \frac{1}{25} \dots\dots(1)$$

$$T_{25} = a + 24d = \frac{1}{20}$$

$$20 \cdot \frac{25}{2} \left[ a + \frac{1}{20} \right] = 13 \Rightarrow a = \frac{1}{500}$$

$$\text{also, } 20S_{25} = 20 \cdot \frac{25}{2} [2a + 24d] = 13 \Rightarrow d = \frac{1}{500}$$

$$\text{from (1) } \frac{1}{500} + \frac{m-1}{500} = \frac{1}{25} \Rightarrow m = 20$$

Now,

$$5m \sum_{r=m}^{2m} T_r = 100 \sum_{r=20}^{40} T_r = 126$$

12. If the image of the point (4, 4, 3) in the line

$$\frac{x-1}{2} = \frac{y-2}{1} = \frac{z-1}{3} \text{ is } (\alpha, \beta, \gamma), \text{ then } \alpha + \beta + \gamma \text{ is}$$

equal to


(1) 9

(2) 12

(3) 8

(4) 7

Ans. (1)

Sol. 

$$\overline{PQ} \perp (2\hat{i} + \hat{j} + 3\hat{k})$$

$$\Rightarrow 2(2\lambda - 3) + 1(\lambda - 2) + 3(3\lambda - 2) = 0$$

$$\Rightarrow 14\lambda - 14 = 0, \lambda = 1$$

$$\text{So, } Q(3, 3, 4)$$

$$\text{Let image in } R(\alpha, \beta, \gamma)$$

$$\frac{\alpha+4}{2} = 3, \frac{\beta+4}{2} = 3, \frac{\gamma+3}{2} = 4$$

$$(\alpha, \beta, \gamma) = (2, 2, 5)$$

$$\Rightarrow \alpha + \beta + \gamma = 9$$

13. If  $\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{96x^2 \cos^2 x}{(1+e^x)} dx = \pi(\alpha\pi^2 + \beta), \alpha, \beta \in \mathbb{Z}$ , then

$$(\alpha + \beta)^2 \text{ equals :}$$

(1) 144

(2) 196

(3) 100

(4) 64

Ans. (3)

Sol.  $\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{96x^2 \cos^2 x}{(1+e^x)} dx$  (Apply King Property)



$$\int_0^{\frac{\pi}{2}} 96x^2 \cos^2 x = 48 \int_0^{\frac{\pi}{2}} x^2 (1 + \cos 2x) dx$$

$$48 \left[ \left( \frac{x^3}{3} \right)_0^{\frac{\pi}{2}} + \int_0^{\frac{\pi}{2}} x^2 \cos 2x dx \right]$$

$\Rightarrow$  On solving  $\pi(2\pi^2 - 12)$

$\Rightarrow \alpha = 2, \beta = -12$

$\Rightarrow (\alpha + \beta)^2 = 100$

14. The sum of all local minimum values of the

$$\text{function } f(x) = \begin{cases} 1 - 2x, & x < -1 \\ \frac{1}{3}(7 + 2|x|), & -1 \leq x \leq 2 \\ \frac{11}{18}(x - 4)(x - 5), & x > 2 \end{cases}$$

(1)  $\frac{171}{72}$


(2)  $\frac{131}{72}$

(3)  $\frac{157}{72}$

(4)  $\frac{167}{72}$

Ans. (3)

Sol.  $f(x) = \begin{cases} 1 - 2x, & x < -1 \\ \frac{1}{3}(7 - 2x), & -1 \leq x \leq 2 \\ \frac{1}{3}(7 + 2x), & 0 \leq x < 2 \\ \frac{11}{18}(x - 4)(x - 5), & x > 2 \end{cases}$



$\therefore$  Local minimum values at A & B

$$\frac{7}{3} - \frac{11}{72}$$

$$\Rightarrow \frac{168 - 11}{72} \Rightarrow \frac{157}{72}$$

15. The sum, of the squares of all the roots of the equation  $x^2 + |2x - 3| - 4 = 0$ , is :

(1)  $3(3 - \sqrt{2})$

(2)  $6(3 - \sqrt{2})$

(3)  $6(2 - \sqrt{2})$

(4)  $3(2 - \sqrt{2})$

Ans. (3)

Sol.  $x^2 + |2x - 3| - 4 = 0$

Case I :  $x \geq \frac{3}{2}$

$$x^2 + 2x - 3 - 4 = 0$$

$$x^2 + 2x - 7 = 0$$

$$x = 2\sqrt{2} - 1$$

Case II :  $x < \frac{3}{2}$

$$x^2 + 3 - 2x - 4 = 0$$

$$x^2 - 2x - 1 = 0$$

$$x = 1 - \sqrt{2}$$

$$\begin{aligned} \text{Sum of squares} &= (2\sqrt{2} - 1)^2 + (1 - \sqrt{2})^2 \\ &= 8 - 4\sqrt{2} + 1 + 1 - 2\sqrt{2} + 2 \\ &= 6(2 - \sqrt{2}) \quad \therefore (3) \end{aligned}$$

16. Let for some function  $y = f(x)$ ,  $\int_0^x t f(t) dt = x^2 f(x)$ ,

$x > 0$  and  $f(2) = 3$ . Then  $f(6)$  is equal to :

(1) 1

(2) 2

(3) 6

(4) 3

Ans. (1)

Sol.  $\int_0^x t f(t) dt = x^2 + (x), x > 0$

Diff. both side w.r. to  $x$

$$x f(x) = x^2 f'(x) + 2x f(x)$$

$$-x f(x) = x^2 f''(x)$$

$$\int \frac{f'(x)}{f(x)} dx = \int \frac{-1}{x} dx$$

$$\log df(x) = -\log x + \log c$$

$$f(x) = \frac{c}{x}$$

$$f(2) = 3 \Rightarrow 3 = \frac{c}{2} \Rightarrow c = 6$$

$$f(x) = \frac{6}{x}$$

$$f(6) = 1 \quad \therefore (1)$$



17. Let  ${}^nC_{r-1} = 28$ ,  ${}^nC_r = 56$  and  ${}^nC_{r+1} = 70$ . Let  $A(4\cos t, 4\sin t)$ ,  $B(2\sin t, -2\cos t)$  and  $C(3r-n, r^2-n-1)$  be the vertices of a triangle ABC, where  $t$  is a parameter. If  $(3x-1)^2 + (3y)^2 = \alpha$ , is the locus of the centroid of triangle ABC, then  $\alpha$  equals :

- (1) 20 (2) 8  
(3) 6 (4) 18

Ans. (1)

Sol.  ${}^nC_{r-1} = 28$ ,  ${}^nC_r = 56$

$$\frac{{}^nC_{r-1}}{{}^nC_r} = \frac{28}{56}$$

$$\frac{\frac{n!}{(r-1)!(n-r+1)!}}{\frac{n!}{r!(n-r)!}} = \frac{1}{2}$$

$$\frac{r}{(n-r+1)} = \frac{1}{2}$$

$$3r = n+1 \quad \text{---(i)}$$

$$\frac{{}^nC_r}{{}^nC_{r+1}} = \frac{56}{70}$$

$$\frac{(r+1)}{(n-r)} = \frac{56}{70} \Rightarrow 9r = 4n-5 \quad \text{---(ii)}$$

By (i) & (ii)

$$(r=3), (n=8)$$

$$A(4\cos t, 4\sin t) \quad B(2\sin t, -2\cos t) \quad C(3r-n, r^2-n-1)$$

$$A(4\cos t, 4\sin t) \quad B(2\sin t, -2\cos t) \quad C(1, 0)$$

$$(3x-1)^2 + (3y)^2 = (4\cos t + 2\sin t)^2 + (4\sin t - \cos t)^2$$

$$(3x-1)^2 + (3y)^2 = 20 \quad \therefore \text{option (1)}$$

18. Let O be the origin, the point A be  $z_1 = \sqrt{3} + 2\sqrt{2}i$ , the point B( $z_2$ ) be such that

$$\sqrt{3}|z_2| = |z_1| \text{ and } \arg(z_2) = \arg(z_1) + \frac{\pi}{6}. \text{ Then}$$

$$(1) \text{ area of triangle ABO is } \frac{11}{\sqrt{3}}$$

$$(2) \text{ ABO is a scalene triangle}$$

$$(3) \text{ area of triangle ABO is } \frac{11}{4}$$

$$(4) \text{ ABO is an obtuse angled isosceles triangle}$$

Ans. (4)

$$\text{Sol. } z_1 = \sqrt{3} + 2\sqrt{2}i \text{ \& } \frac{|z_2|}{|z_1|} = \frac{1}{\sqrt{3}}$$

$$\text{given } \arg\left(\frac{z_2}{z_1}\right) = \frac{\pi}{6}$$

$$z_2 = \frac{|z_2|}{|z_1|} \cdot z_1 \cdot e^{i\left(\frac{\pi}{6}\right)}$$

$$z_2 = \frac{1}{\sqrt{3}} \cdot \frac{(\sqrt{3} + 2\sqrt{2}i)(\sqrt{3} + i)}{2}$$

$$z_2 = \frac{(3 - 2\sqrt{2}) + i(2\sqrt{6} + \sqrt{3})}{2\sqrt{3}}$$

Now,

$$z_1 - z_2 = \frac{(3 + 2\sqrt{2}) + i(2\sqrt{6} - \sqrt{3})}{2\sqrt{3}}$$

$$|z_1 - z_2| = |z_2| \Rightarrow \Delta ABO \text{ is isosceles with angles } \frac{\pi}{6}, \frac{\pi}{6} \text{ \& } \frac{2\pi}{3}$$

19. Three defective oranges are accidentally mixed with seven good ones and on looking at them, it is not possible to differentiate between them. Two oranges are drawn at random from the lot. If  $x$  denote the number of defective oranges, then the variance of  $x$  is :

- (1) 28/75 (2) 14/25  
(3) 26/75 (4) 18/25

Ans. (1)

Sol. 10 oranges  $\begin{cases} 7 \text{ good} \\ 3 \text{ defected} \end{cases}$

Probability distribution

$x_i$	$p_i$
$x=0$	$\frac{{}^7C_2}{{}^{10}C_2} = \frac{42}{90}$
$x=1$	$\frac{{}^7C_1 \times {}^3C_1}{{}^{10}C_2} = \frac{42}{90}$
$x=2$	$\frac{{}^3C_2}{{}^{10}C_2} = \frac{6}{90}$

Now,

$$\mu = \sum x_i p_i = \frac{42}{90} + \frac{12}{90} = \frac{54}{90}$$

$$\sigma^2 = \sum p_i x_i^2 - \mu^2 = \frac{42}{90} + \frac{24}{90} - \left(\frac{54}{90}\right)^2$$

$$\Rightarrow \frac{66}{90} - \left(\frac{54}{90}\right)^2$$


$$\sigma^2 \Rightarrow \frac{28}{75} \therefore (1)$$



20. The area (in sq. units) of the region  
 $\{(x, y): 0 \leq y \leq 2|x| + 1, 0 \leq y \leq x^2 + 1, |x| \leq 3\}$   
 is

- (1)  $\frac{80}{3}$  (2)  $\frac{64}{3}$   
 (3)  $\frac{17}{3}$  (4)  $\frac{32}{3}$

Ans. (2)



Sol.

$$\text{Area} = 2 \left[ \int_0^3 (x^2 + 1) dx + \int_2^3 (2x + 1) dx \right]$$

$$\Rightarrow \frac{64}{3} \quad \therefore (2)$$

### SECTION-B

21. Let  $M$  denote the set of all real matrices of order  $3 \times 3$  and let  $S = \{-3, -2, -1, 1, 2\}$ . Let  
 $S_1 = \{A = [a_{ij}] \in M : A = A^T \text{ and } a_{ij} \in S, \forall i, j\}$   
 $S_2 = \{A = [a_{ij}] \in M : A = -A^T \text{ and } a_{ij} \in S, \forall i, j\}$   
 $S_3 = \{A = [a_{ij}] \in M : a_{11} + a_{22} + a_{33} = 0 \text{ and } a_{ij} \in S, \forall i, j\}$   
 If  $n(S_1 \cup S_2 \cup S_3) = 125\alpha$ , then  $\alpha$  equals.

Ans. (1613)

Sol. 
$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

No. of elements in  $S_1 : A = A^T \Rightarrow 5^3 \times 5^3$

No. of elements in  $S_2 : A = -A^T \Rightarrow 0$

since no zero in  $S_2$

No. of elements in  $S_3 \Rightarrow$

$$a_{11} + a_{22} + a_{33} = 0 \Rightarrow (1, 2, -3) \Rightarrow 31$$

or

$$(1, 1, -2) \Rightarrow 3$$

or

$$(-1, -1, 2) \Rightarrow 3$$

$$\left. \begin{array}{l} 31 \\ 3 \\ 3 \end{array} \right\} \Rightarrow 12 \times 5^6$$

$$n(S_1 \cap S_3) = 12 \times 5^3$$

$$n(S_1 \cup S_2 \cup S_3) = 5^6(1+12) - 12 \times 5^3$$

$$\Rightarrow 5^3 \times [13 \times 5^3 - 12] = 125\alpha$$

$$\alpha = 1613$$

22. If  $\alpha = 1 + \sum_{r=1}^6 (-3)^{r-1} {}^{12}C_{2r-1}$ , then the distance of the point  $(12, \sqrt{3})$  from the line  $\alpha x - \sqrt{3}y + 1 = 0$  is

Ans. (5)

Sol. 
$$\alpha = 1 + \sum_{r=1}^6 (-1)^{r-1} {}^{12}C_{2r-1} 3^{r-1}$$

$$\alpha = 1 + \sum_{r=1}^6 {}^{12}C_{2r-1} \frac{(\sqrt{3}i)^{2r-1}}{\sqrt{3}i} \quad i = \text{iota, let } \sqrt{3}i = x$$

$$\alpha = 1 + \frac{1}{\sqrt{3}i} \left( {}^{12}C_1 x + {}^{12}C_3 x^3 + \dots + {}^{12}C_{11} x^{11} \right)$$

$$= 1 + \frac{1}{\sqrt{3}i} \left( \frac{(1 + \sqrt{3}i)^{12} - (1 - \sqrt{3}i)^{12}}{2} \right)$$

$$= 1 + \frac{1}{\sqrt{3}i} \left( \frac{(-2\omega^2)^{12} - (2\omega)^{12}}{2} \right) = 1$$

so distance of  $(12, \sqrt{3})$  from  $x - \sqrt{3}y + 1 = 0$  is

$$\frac{12 - 3 + 1}{2} = 5$$

23. Let  $\vec{a} = \hat{i} + \hat{j} + \hat{k}$ ,  $\vec{b} = 2\hat{i} + 2\hat{j} + \hat{k}$  and  $\vec{d} = \vec{a} \times \vec{b}$ .  
 If  $\vec{c}$  is a vector such that  $\vec{a} \cdot \vec{c} = |\vec{c}|$ ,  $|\vec{c} - 2\vec{a}|^2 = 8$   
 and the angle between  $\vec{d}$  and  $\vec{c}$  is  $\frac{\pi}{4}$ , then

$$|10 - 3\vec{b} \cdot \vec{c}| + |\vec{d} \times \vec{c}|^2 \text{ is equal to } \dots$$

Ans. (6)

Sol. 
$$\vec{a} = \hat{i} + \hat{j} + \hat{k}$$

$$\vec{b} = 2\hat{i} + 2\hat{j} + \hat{k}$$

$$\vec{d} = \vec{a} \times \vec{b}$$

$$= -\hat{i} + \hat{j}$$

$$|\vec{c} - 2\vec{a}|^2 = 8$$

$$|\vec{c}|^2 + 4|\vec{a}|^2 - 4(\vec{a} \cdot \vec{c}) = 8$$

$$c^2 + 12 - 4c = 8$$

$$c^2 - 4c + 4 = 0$$

$$|c| = 2$$

$$\vec{d} = \vec{a} \times \vec{b}$$

$$\vec{d} \times \vec{c} = (\vec{a} \times \vec{b}) \times \vec{c}$$

$$\left( |\vec{d}| |\vec{c}| \sin \frac{\pi}{4} \right)^2 = ((\vec{a} \cdot \vec{c}) \cdot \vec{b} - (\vec{b} \cdot \vec{c}) \cdot \vec{a})^2$$

$$4 = 4b^2 + (b \cdot c)^2 2(a)^2 - 2(b \cdot c)(a \cdot b)$$



Let  $b.c = x$

$$4 = 36 + 3x^2 - 20x$$

$$3x^2 - 20x + 32 = 0$$

$$3x^2 - 12x - 8x + 32 = 0$$

$$x = \frac{8}{3}, 4$$

$$b.c = \frac{8}{3}, 4$$

$$b.c = \frac{8}{3}$$

Now  $|10 - 3b.c| + |d \times c|^2$

$$|10 - 8| + (2)^2$$

$$\Rightarrow 6 \text{ Ans.}$$

24. Let


$$f(x) = \begin{cases} 3x, & x < 0 \\ \min\{1+x+[x], x+2[x]\}, & 0 \leq x \leq 2 \\ 5, & x > 2 \end{cases}$$

where  $[.]$  denotes greatest integer function. If  $\alpha$  and  $\beta$  are the number of points, where  $f$  is not continuous and is not differentiable, respectively, then  $\alpha + \beta$  equals.....

Ans. (5)

Sol.  $f(x) = \begin{cases} 3x & ; x < 0 \\ \min\{1+x, x\} & ; 0 \leq x < 1 \\ \min\{2+x, x+2\} & ; 1 \leq x < 2 \\ 5 & ; x > 2 \end{cases}$

$$f(x) = \begin{cases} 3x & ; x < 0 \\ x & ; 0 \leq x < 1 \\ x+2 & ; 1 \leq x < 2 \\ 5 & ; x > 2 \end{cases}$$




Not continuous at  $x \in \{0, 1, 2\} \Rightarrow \alpha = 3$

Not diff. at  $x \in \{0, 1, 2\} \Rightarrow \beta = 3$

$$\alpha + \beta = 6$$

25. Let  $E_1 : \frac{x^2}{9} + \frac{y^2}{4} = 1$  be an ellipse. Ellipses  $E_i$ 's are constructed such that their centres and eccentricities are same as that of  $E_1$ , and the length of minor axis of  $E_i$  is the length of major axis of  $E_{i+1}$  ( $i \geq 1$ ). If  $A_i$  is the area of the ellipse  $E_i$ , then  $\frac{5}{\pi} \left( \sum_{i=1}^{\infty} A_i \right)$ , is equal to .....

Ans. (54)



Sol.

$$E_1 : \frac{x^2}{9} + \frac{y^2}{4} = 1 \Rightarrow e = \sqrt{1 - \frac{4}{9}} = \frac{\sqrt{5}}{3}$$

$$E_2 : \frac{x^2}{a^2} + \frac{y^2}{4} = 1$$

$$e = \frac{\sqrt{5}}{3} = \sqrt{1 - \frac{a^2}{4}} \Rightarrow \frac{5}{9} = 1 - \frac{a^2}{4}$$

$$a^2 = \frac{16}{9}$$

$$E_2 : \frac{x^2}{\frac{16}{9}} + \frac{y^2}{4} = 1$$

$$E_3 : \frac{x^2}{\frac{16}{9}} + \frac{y^2}{b^2} = 1$$

$$e = \frac{\sqrt{5}}{3} = \sqrt{1 - \frac{b^2}{\frac{16}{9}}} \Rightarrow b^2 = \frac{64}{81}$$

$$E_3 : \frac{x^2}{\frac{16}{9}} + \frac{y^2}{\frac{64}{81}} = 1$$

$$A_1 = \pi \times 3 \times 2 \Rightarrow 6\pi$$

$$A_2 = \pi \times \frac{4}{3} \times 2 = \frac{8\pi}{3}$$

$$A_3 = \pi \times \frac{4}{3} \times \frac{8}{9} = \frac{32\pi}{81}$$

$$\sum_{i=1}^{\infty} A_i = 6\pi + \frac{8\pi}{3} + \frac{32\pi}{81} + \dots \Rightarrow \frac{6\pi}{1 - \frac{4}{9}} \Rightarrow \frac{54\pi}{5}$$


$$\therefore \frac{5}{\pi} \sum_{i=1}^{\infty} A_i \Rightarrow \frac{5}{\pi} \times \frac{54\pi}{5} = 54$$



## PHYSICS

### SECTION-A

26. Two capacitors  $C_1$  and  $C_2$  are connected in parallel to a battery. Charge-time graph is shown below for the two capacitors. The energy stored with them are  $U_1$  and  $U_2$ , respectively. Which of the given statements is true ?



- (1)  $C_1 > C_2$ ,  $U_1 > U_2$       (2)  $C_2 > C_1$ ,  $U_2 < U_1$   
 (3)  $C_1 > C_2$ ,  $U_1 < U_2$       (4)  $C_2 > C_1$ ,  $U_2 > U_1$

**Ans. (4)**

**Sol.** potential difference,

$v \rightarrow$  same

$$U = \frac{1}{2} C v^2$$

as  $q_1 < q_2$

$\therefore C_1 < C_2$

&  $U_1 < U_2$

27. In the experiment for measurement of viscosity ' $\eta$ ' of given liquid with a ball having radius  $R$ , consider following statements.

- A. Graph between terminal velocity  $V$  and  $R$  will be a parabola  
 B. The terminal velocities of different diameter balls are constant for a given liquid.  
 C. Measurement of terminal velocity is dependent on the temperature.  
 D. This experiment can be utilized to assess the density of a given liquid.  
 E. If balls are dropped with some initial speed, the value of  $\eta$  will change.

Choose the correct answer from the options given below:

- (1) B, D and E only  
 (2) A, C and D only  
 (3) C, D and E only  
 (4) A, B and E only

**Ans. (2)**

**Sol.**  $V_T = \frac{2}{9} R^2 \frac{g}{\eta} (d - \rho)$

28. Consider following statements:

- A. Surface tension arises due to extra energy of the molecules at the interior as compared to the molecules at the surface, of a liquid.  
 B. As the temperature of liquid rises, the coefficient of viscosity increases.  
 C. As the temperature of gas increases, the coefficient of viscosity increases.  
 D. The onset of turbulence is determined by Reynold's number.  
 E. In a steady flow two stream lines never intersect.

Choose the correct answer from the options given below :

- (1) A, D, E only  
 (2) C, D, E only  
 (3) B, C, D only  
 (4) A, B, C only

**Ans. (2)**

29. Three infinitely long wires with linear charge density  $\lambda$  are placed along the x-axis, y-axis and z-axis respectively. Which of the following denotes an equipotential surface ?

- (1)  $xy + yz + zx = \text{constant}$   
 (2)  $(x + y)(y + z)(z + x) = \text{constant}$   
 (3)  $(x^2 + y^2)(y^2 + z^2)(z^2 + x^2) = \text{constant}$   
 (4)  $xyz = \text{constant}$



Ans. (3)

Sol.  $v = -\int \vec{E} \cdot d\vec{r} = \int \frac{2k\lambda}{r} dr = 2k\lambda \ln r + c$

Net potential due to all wire

$$v = 2k\lambda \ln \sqrt{x^2 + y^2} + 2k\lambda \ln \sqrt{y^2 + z^2} + 2k\lambda \ln \sqrt{z^2 + x^2} + c$$


for  $v = c$

$$\sqrt{(x^2 + y^2)(y^2 + z^2)(z^2 + x^2)} = c$$

$$\therefore (x^2 + y^2)(y^2 + z^2)(z^2 + x^2) = c$$

where  $c = \text{constant}$


30. A hemispherical vessel is completely filled with a liquid of refractive index  $\mu$ . A small coin is kept at the lowest point (O) of the vessel as shown in figure. The minimum value of the refractive index of the liquid so that a person can see the coin from point E (at the level of the vessel) is \_\_\_\_\_.



- (1)  $\sqrt{3}$  (2)  $\frac{3}{2}$   
(3)  $\sqrt{2}$  (4)  $\frac{\sqrt{3}}{2}$

Ans. (3)

Sol.



$$\sin c = \frac{1}{\mu}$$

for  $\mu \rightarrow \text{least}$ ,  $c \rightarrow \text{maximum}$


$$\theta = c = 45$$

$$\mu = \frac{1}{\sin 45} = \sqrt{2}$$

31. Consider a long thin conducting wire carrying a uniform current  $I$ . A particle having mass " $M$ " and charge " $q$ " is released at a distance " $a$ " from the wire with a speed  $v_0$  along the direction of current in the wire. The particle gets attracted to the wire due to magnetic force. The particle turns round when it is at distance  $x$  from the wire. The value of  $x$  is [ $\mu_0$  is vacuum permeability]

- (1)  $a \left[ 1 - \frac{mv_0}{2q\mu_0 I} \right]$  (2)  $\frac{a}{2}$   
(3)  $a \left[ 1 - \frac{mv_0}{q\mu_0 I} \right]$  (4)  $ae^{\frac{-4\pi mv_0}{q\mu_0 I}}$

Ans. (4)



Sol.

$A \rightarrow B$

$$\vec{V} = -v_x \hat{i} + v_y \hat{j}$$

$$\vec{B} = \frac{\mu_0 I}{2\pi r} (-\hat{k})$$

$$\vec{F} = q(\vec{V} \times \vec{B}) = \frac{\mu_0 I q}{2\pi r} [-v_x \hat{j} - v_y \hat{i}]$$

$$a_x = -\frac{\mu_0 I q}{2\pi m} \cdot \frac{v_y}{r}$$

$$a_y = -\frac{\mu_0 I q}{2\pi m} \cdot \frac{v_x}{r}$$

$$\frac{v_x dv_x}{dr} = -\frac{\mu_0 I q}{2\pi m} \cdot \frac{v_y}{r}$$

$$\frac{v_x dv_x}{v_y} = -\frac{\mu_0 I q}{2\pi m} \cdot \frac{dr}{r}$$

$$\int_0^{v_0} \frac{v_x dv_x}{\sqrt{v_0^2 - v_x^2}} = -\frac{\mu_0 I q}{2\pi m} \int_a^{x_1} \frac{dr}{r}$$

$$\text{Let, } z^2 = v_0^2 - v_x^2$$

$$2z dz = -2v_x dv_x$$



$$\frac{v_x dv_x}{\sqrt{v_0^2 - v_x^2}} = \frac{-z dz}{z} = -dz$$

then integral becomes

$$-\int_{v_0}^0 dz = -\frac{\mu_0 I q}{2\pi m} \ln \frac{x_1}{a}$$

$$v_0 = -\frac{\mu_0 I q}{2\pi m} \ln \frac{x_1}{a}$$

$$x_1 = a e^{\frac{2\pi m v_0}{\mu_0 I q}} \dots\dots(1)$$

For B  $\rightarrow$  C

$$\vec{v} = -v_x \hat{i} - v_y \hat{j}$$

$$\vec{B} = \frac{\mu_0 I}{2\pi r} (-\hat{k})$$

$$\vec{F} = q(\vec{v} \times \vec{B}) = \frac{\mu_0 I q}{2\pi r} (-v_x \hat{j} + v_y \hat{i})$$

$$a_x = +\frac{\mu_0 I q}{2\pi m} \frac{v_y}{r} \quad a_y = -\frac{\mu_0 I q}{2\pi m} \cdot \frac{v_x}{r}$$

$$\frac{v_x dv_x}{dr} = \frac{\mu_0 I q}{2\pi m} \frac{v_y}{r}$$

$$\int_{v_0}^0 \frac{v_x dv_x}{\sqrt{v_0^2 - v_x^2}} = \frac{\mu_0 I q}{2\pi m} \int_{x_1}^x \frac{dr}{r}$$

$$\frac{\mu_0 I q}{2\pi m} \ln \frac{x}{x_1} = -\int_0^{v_0} dz = -v_0$$

$$x = x_1 e^{\frac{2\pi m v_0}{\mu_0 I q}} \dots\dots(2)$$

From equation 1 and 2

$$X = a e^{\frac{4\pi m v_0}{\mu_0 I q}}$$

32. A Carnot engine (E) is working between two temperatures 473K and 273K. In a new system two engines – engine  $E_1$  works between 473K to 373K and engine  $E_2$  works between 373K to 273K. If  $\eta_{12}$ ,  $\eta_1$  and  $\eta_2$  are the efficiencies of the engines E,  $E_1$  and  $E_2$ , respectively, then

- (1)  $\eta_{12} < \eta_1 + \eta_2$  (2)  $\eta_{12} = \eta_1 \eta_2$   
 (3)  $\eta_{12} = \eta_1 + \eta_2$  (4)  $\eta_{12} \geq \eta_1 + \eta_2$

Ans. (1)

Sol.  $\eta_{12} = 1 - \frac{273}{473} = \frac{200}{473} = 0.423$

$$\eta_1 = 1 - \frac{373}{473} = \frac{100}{473} = 0.211$$

$$\eta_2 = 1 - \frac{273}{373} = \frac{100}{373} = 0.268$$

33. Given below are two statements: one is labelled as **Assertion A** and the other is labelled as **Reason R**  
**Assertion A:** A sound wave has higher speed in solids than gases.

**Reason R:** Gases have higher value of Bulk modulus than solids.


In the light of the above statements, choose the **correct** answer from the options given below

- (1) Both **A** and **R** are true and **R** is the correct explanation of **A**  
 (2) **A** is false but **R** is true  
 (3) Both **A** and **R** are true but **R** is **NOT** the correct explanation of **A**  
 (4) **A** is true but **R** is false.

Ans. (4)

- Sol. Solids have higher value of bulk modulus than gases.

34. For a particular ideal gas which of the following graphs represents the variation of mean squared velocity of the gas molecules with temperature ?



Ans. (1)

Sol.  $V_{rms} = \sqrt{\frac{3RT}{M}}$

$$V_{rms}^2 = 3RT/M$$


Hence we can conclude that  $V_{rms}^2$  is directly proportional to temperature

$$y = m x$$

$\Rightarrow$  Graph will be straight line




35. A bead of mass 'm' slides without friction on the wall of a vertical circular hoop of radius 'R' as shown in figure. The bead moves under the combined action of gravity and a massless spring (k) attached to the bottom of the hoop. The equilibrium length of the spring is 'R'. If the bead is released from top of the hoop with (negligible) zero initial speed, velocity of bead, when the length of spring becomes 'R', would be (spring constant is 'k', g is acceleration due to gravity)



- (1)  $2\sqrt{gR + \frac{kR^2}{m}}$  (2)  $\sqrt{2Rg + \frac{4kR^2}{m}}$   
 (3)  $\sqrt{2Rg + \frac{kR^2}{m}}$  (4)  $\sqrt{3Rg + \frac{kR^2}{m}}$

Ans. (4)



Sol.

Work energy theorem

$$Mg(R + R\cos 60) + \frac{1}{2}k(R^2 - 0^2) = \frac{1}{2}mv^2$$

$$Mg \frac{3R}{2} + \frac{KR^2}{2} = \frac{1}{2}mv^2$$

$$V = \sqrt{3gR + \frac{KR^2}{m}}$$

36. Given below are two statements: one is labelled as **Assertion A** and the other is labelled as **Reason R**  
**Assertion A:** In a central force field, the work done is independent of the path chosen  
**Reason R:** Every force encountered in mechanics does not have an associated potential energy.

In the light of the above statements, choose the **most appropriate** answer from the options given below

- (1) **A** is true but **R** is false  
 (2) Both **A** and **R** are true but **R** is **NOT** the correct explanation of **A**  
 (3) Both **A** and **R** are true and **R** is the correct explanation of **A**  
 (4) **A** is false but **R** is true

Ans. (2)

Sol. Both statement are correct but Reason is not the correct explanation of Assertion.


37. Choose the correct nuclear process from the below options

[p: proton, n: neutron,  $e^-$ : electron,  $e^+$ : positron,  $\nu$ : neutrino,  $\bar{\nu}$ : antineutrino]


- (1)  $n \rightarrow p + e^- + \bar{\nu}$  (2)  $n \rightarrow p + e^- + \nu$   
 (3)  $n \rightarrow p + e^+ + \bar{\nu}$  (4)  $n \rightarrow p + e^+ + \nu$

Ans. (1)

Sol. Theoretical equation for  $\beta^-$  decay




38. Which of the following circuits has the same output as that of the given circuit?



- (1)   
 (2)   
 (3)   
 (4)

Ans. (1)





$$P = A \cdot \bar{B}$$


$$Q = A \cdot B$$

$$Y = \overline{P+Q} = \overline{A \cdot \bar{B} + A \cdot B}$$

$$= \overline{A \cdot (B + \bar{B})} = \overline{A \cdot 1}$$



$$Y = \bar{A}$$

- 39.** Find the equivalent resistance between two ends of the following circuit.



- (1)  $r$  (2)  $\frac{r}{6}$   
 (3)  $\frac{r}{9}$  (4)  $\frac{r}{3}$

**Ans. (3)**



All are in parallel

$$R_{eq} = \frac{r/3}{3} = r/9$$

- 40.** A wire of resistance  $R$  is bent into an equilateral triangle and an identical wire is bent into a square. The ratio of resistance between the two end points of an edge of the triangle to that of the square is


- (1)  $9/8$  (2)  $8/9$   
 (3)  $27/32$  (4)  $32/27$

**Ans. (4)**

**Sol.**  $R = \frac{\ell}{A}$

So,  $R \propto \ell$

Side length of triangle is  $1/3$  of total length.



$$(R_{eq})_1 = \frac{2r/3 \times r/3}{2r/3 + r/3} \quad (R_{eq})_2 = \frac{3r/4 \times r/4}{3r/4 + r/4}$$

$$(R_{eq})_1 = 2r/9 \quad (R_{eq})_2 = 3r/16$$

$$\frac{(R_{eq})_1}{(R_{eq})_2} = \frac{2r/9}{3r/16} = \frac{32}{27}$$

- 41.** Due to presence of an em-wave whose electric component is given by  $E = 100 \sin(\omega t - kx) \text{ NC}^{-1}$ , a cylinder of length 200 cm holds certain amount of em-energy inside it. If another cylinder of same length but half diameter than previous one holds same amount of em-energy, the magnitude of the electric field of the corresponding em-wave should be modified as

- (1)  $25 \sin(\omega t - kx) \text{ NC}^{-1}$   
 (2)  $200 \sin(\omega t - kx) \text{ NC}^{-1}$   
 (3)  $400 \sin(\omega t - kx) \text{ NC}^{-1}$   
 (4)  $50 \sin(\omega t - kx) \text{ NC}^{-1}$

**Ans. (2)**

**Sol.** Energy density  $= \frac{1}{2} \epsilon_0 E^2 \times c$

$$\text{Energy} = \frac{1}{2} \epsilon_0 E^2 \times c \times \text{volume}$$

$$(\text{Energy})_1 = (\text{Energy})_2 \quad (\text{Given})$$

$$\frac{1}{2} \epsilon_0 E_1^2 c \pi R_1^2 \times L_1 = \frac{1}{2} \epsilon_0 E_2^2 c \pi R_2^2 \times L_2$$

$$E_1^2 R_1^2 = E_2^2 R_2^2$$


$$E_1 R_1 = E_2 R_2$$

$$100 \times R_1 = E_2 \times \frac{R_1}{2}$$

$$E_2 = 200 \text{ N/C}$$




42. A particle of mass 'm' and charge 'q' is fastened to one end 'A' of a massless string having equilibrium length  $\ell$ , whose other end is fixed at point 'O'. The whole system is placed on a frictionless horizontal plane and is initially at rest. If uniform electric field is switched on along the direction as shown in figure, then the speed of the particle when it crosses the x-axis is



- (1)  $\sqrt{\frac{2qE\ell}{m}}$  (2)  $\sqrt{\frac{qE\ell}{4m}}$   
 (3)  $\sqrt{\frac{qE\ell}{m}}$  (4)  $\sqrt{\frac{qE\ell}{2m}}$

Ans. (3)

Sol.



$$W_{\text{all}} = \Delta k$$

$$W_e = k_f - k_i$$

$$qE \frac{\ell}{2} = \frac{1}{2}mv^2 - 0$$

$$v = \sqrt{\frac{qE\ell}{m}}$$

43. A proton of mass ' $m_p$ ' has same energy as that of a photon of wavelength ' $\lambda$ '. If the proton is moving at non-relativistic speed, then ratio of its de Broglie wavelength to the wavelength of photon is.

- (1)  $\frac{1}{c} \sqrt{\frac{2E}{m_p}}$  (2)  $\frac{1}{c} \sqrt{\frac{E}{m_p}}$   
 (3)  $\frac{1}{c} \sqrt{\frac{E}{2m_p}}$  (4)  $\frac{1}{2c} \sqrt{\frac{E}{m_p}}$

Ans. (3)

- Sol. E is missing in the question but considering E as energy, the solution will be


$$E_{\text{photon}} = \frac{hc}{\lambda} = E ; E_{\text{proton}} = \frac{1}{2}m_p v^2 = E$$

$$\frac{\lambda_{\text{proton}}}{\lambda_{\text{photon}}} = \frac{h/p}{hc/E} = \frac{h/\sqrt{2m_p E}}{hc/E}$$

$$= \frac{E}{c\sqrt{2m_p E}}$$

$$\frac{\lambda_{\text{proton}}}{\lambda_{\text{photon}}} = \frac{1}{c} \sqrt{\frac{E}{2m_p}}$$

44. The centre of mass of a thin rectangular plate (fig - x) with sides of length a and b, whose mass per unit area ( $\sigma$ ) varies as  $\sigma = \frac{\sigma_0 x}{ab}$  (where  $\sigma_0$  is a constant), would be \_\_\_\_\_




- (1)  $\left(\frac{2}{3}a, \frac{b}{2}\right)$  (2)  $\left(\frac{2}{3}a, \frac{2}{3}b\right)$   
 (3)  $\left(\frac{a}{2}, \frac{b}{2}\right)$  (4)  $\left(\frac{1}{3}a, \frac{b}{2}\right)$

Ans. (1)

- Sol.  $\sigma$  is constant in y-direction

$$\text{So, } y_{\text{cm}} = b/2$$



$$x_{\text{cm}} = \frac{\int_0^a x dm}{\int_0^a dm}$$



$$\begin{aligned}
&= \frac{\int_0^a x \sigma_x dA}{\int_0^a \sigma_x dA} \\
&= \frac{\int_0^a x \frac{\sigma_0 x}{ab} b dx}{\int_0^a \frac{\sigma_0 x}{ab} b dx} \\
&= \frac{\int_0^a x^2 dx}{\int_0^a x dx} \\
&= \frac{\left( \frac{x^3}{3} \right)_0^a}{\left( \frac{x^2}{2} \right)_0^a} = \frac{a^3/3}{a^2/2} \\
&= \frac{2a}{3}
\end{aligned}$$

45. A thin prism  $P_1$  with angle  $4^\circ$  made of glass having refractive index 1.54, is combined with another thin prism  $P_2$  made of glass having refractive index 1.72 to get dispersion without deviation. The angle of the prism  $P_2$  in degrees is

- (1) 4 (2) 3  
(3) 16/3 (4) 1.5

Ans. (2)

Sol.  $\delta_{\text{net}} = 0$

$$(\mu_1 - 1)A_1 - (\mu_2 - 1)A_2 = 0$$

$$(1.54 - 1)4 - (1.72 - 1)A_2 = 0$$


$$A_2 = 3^\circ$$

46. A tiny metallic rectangular sheet has length and breadth of 5 mm and 2.5 mm, respectively. Using a specially designed screw gauge which has pitch of 0.75 mm and 15 divisions in the circular scale, you are asked to find the area of the sheet. In this measurement, the maximum fractional error will

be  $\frac{x}{100}$  where x is \_\_\_\_\_

Ans. (3)

Sol.



Since least count of the instrument can be calculated as

$$\begin{aligned}
\text{Least count} &= \frac{\text{pitch length}}{\text{No. of division on circular scale}} \\
&= \frac{0.75}{15} = 0.05 \text{ mm.}
\end{aligned}$$

Here we are provided  $L = 5 \text{ mm}$  &  $W = 2.5 \text{ mm}$

$L = 5 \text{ mm}$  &  $W = 2.5 \text{ mm}$

$\therefore$  We know that

$$A = L \cdot W$$

For calculating fractional error, we can write

$$\frac{dA}{A} = \frac{dL}{L} + \frac{dW}{W}$$

Here  $dL = dW = 0.05 \text{ mm}$

$$\frac{dA}{A} = \frac{0.05}{5} + \frac{0.05}{2.5}$$


$$\Rightarrow \frac{dA}{A} = \frac{1}{100} + \frac{2}{100} = \frac{3}{100},$$

So,  $x = 3$

47. The moment of inertia of a solid disc rotating along its diameter is 2.5 times higher than the moment of inertia of a ring rotating in similar way. The moment of inertia of a solid sphere which has same radius as the disc and rotating in similar way, is n times higher than the moment of inertia of the given ring. Here,  $n =$  \_\_\_\_\_.

Consider all the bodies have equal masses.

Ans. (4)



Sol.

$$I_1 = \frac{MR_1^2}{4}, I_2 = \frac{MR_2^2}{2}, I_3 = \frac{2MR_1^2}{5}$$

According to problem



$$\frac{I_1}{I_2} = 2.5 \Rightarrow \frac{\frac{MR_1^2}{2}}{\frac{MR_2^2}{2}} = \frac{5}{2} \Rightarrow \frac{R_1^2}{R_2^2} = 5 \dots (1)$$

Now we are provided with information that

$$\frac{I_3}{I_2} = n$$

$$\Rightarrow \frac{\frac{2MR_1^2}{2}}{\frac{MR_2^2}{2}} = n \Rightarrow \frac{4R_1^2}{5R_2^2} = n \dots (2)$$

From Eq', (1) and (2)

$$\Rightarrow n = 4$$

48. In a measurement, it is asked to find modulus of elasticity per unit torque applied on the system. The measured quantity has dimension of  $[M^a L^b T^c]$ . If  $b = 3$ , the value of  $c$  is \_\_\_\_\_

Ans. (4)


Sol.  $\frac{\text{modulus of elasticity}}{\text{Torque}} = \frac{\text{Stress}}{\text{strain} \times \text{torque}}$

$$= \frac{[\text{Force}]}{[\text{Area}] \times [\text{Force} \times \text{length}]}$$

$$= \frac{1}{[\text{Area} \times \text{length}]} = [L^{-3}]$$

49. Two iron solid discs of negligible thickness have radii  $R_1$  and  $R_2$  and moment of inertia  $I_1$  and  $I_2$ , respectively. For  $R_2 = 2R_1$ , the ratio of  $I_1$  and  $I_2$  would be  $1/x$ , where  $x =$  \_\_\_\_\_

Ans. (16)



Given  $R_2 = 2R_1$

$$M_1 = \sigma \times \pi R_1^2 = M_o$$

$$M_2 = \sigma \times \pi R_2^2 = M_o$$

$$M_2 = \sigma \times \pi R_2^2 = \sigma \times \pi [2R_1]^2 = \sigma \times 4\pi R_1^2 = 4M_o$$

$$\frac{I_1}{I_2} = \frac{\frac{M_1 R_1^2}{2}}{\frac{M_2 R_2^2}{2}} = \frac{M_1 R_1^2}{M_2 R_2^2} = \frac{1}{4} \times \frac{1}{4} = \frac{1}{16}$$

50. A double slit interference experiment performed with a light of wavelength 600 nm forms an interference fringe pattern on a screen with  $10^{\text{th}}$  bright fringe having its centre at a distance of 10 mm from the central maximum. Distance of the centre of the same  $10^{\text{th}}$  bright fringe from the central maximum when the source of light is replaced by another source of wavelength 660 nm would be \_\_\_\_\_ mm.

Ans. (11)

- Sol. In case of YDSE the distance of  $n^{\text{th}}$  maxima from central maxima is given by

$$Y = \frac{n\lambda D}{d}$$

Here  $n$ ,  $D$  &  $d$  are same

So,  $y \propto \lambda$

$$\Rightarrow \frac{y_2}{y_1} = \frac{\lambda_2}{\lambda_1} \Rightarrow \frac{y_2}{10 \text{ mm}} = \frac{660 \text{ nm}}{600 \text{ nm}}$$

$$\Rightarrow y_2 = 11 \text{ mm}$$



## CHEMISTRY

### SECTION-A

51. The incorrect decreasing order of atomic radii is :

- (1)  $\text{Mg} > \text{Al} > \text{C} > \text{O}$       (2)  $\text{Al} > \text{B} > \text{N} > \text{F}$   
 (3)  $\text{Be} > \text{Mg} > \text{Al} > \text{Si}$     (4)  $\text{Si} > \text{P} > \text{Cl} > \text{F}$

Ans. (3)

Sol. Correct order of atomic radii :  $\text{Be} < \text{Mg} > \text{Al} > \text{Si}$

52. Given below are two statements :

**Statement I :** In the oxalic acid vs  $\text{KMnO}_4$  (in the presence of dil  $\text{H}_2\text{SO}_4$ ) titration the solution needs to be heated initially to  $60^\circ\text{C}$ , but no heating is required in Ferrous ammonium sulphate (FAS) vs  $\text{KMnO}_4$  titration (in the presence of dil  $\text{H}_2\text{SO}_4$ )

**Statement II :** In oxalic acid vs  $\text{KMnO}_4$  titration, the initial formation of  $\text{MnSO}_4$  takes place at high temperature, which then acts as catalyst for further reaction. In the case of FAS vs  $\text{KMnO}_4$ , heating oxidizes  $\text{Fe}^{2+}$  into  $\text{Fe}^{3+}$  by oxygen of air and error may be introduced in the experiment.

In the light of the above statements, choose the **correct** answer from the options given below :

- (1) Statement I is false but Statement II is true  
 (2) Both Statement I and Statement II are true  
 (3) Statement I is true but Statement II is false  
 (4) Both Statement I and Statement II are false

Ans. (2)

Sol. For the titration : Oxalic acid v/s  $\text{KMnO}_4$



This reaction is slow at room temperature, but becomes fast at  $60^\circ\text{C}$ . Manganese(II) ions catalyse the reaction; thus, the reaction is autocatalytic; once manganese(II) ions are formed, it becomes faster and faster.

The titration of FAS v/s  $\text{KMnO}_4$  do not require heating because at higher temperature the oxidation of  $\text{Fe}^{+2}$  to  $\text{Fe}^{+3}$  by atmospheric  $\text{O}_2$  will be prominent.

53. Match the **List-I** with **List-II**

List-I (Redox Reaction)		List-II (Type of Redox Reaction)	
A	$\text{CH}_4(\text{g}) + 2\text{O}_2(\text{g}) \xrightarrow{\Delta} \text{CO}_2(\text{g}) + 2\text{H}_2\text{O}(\text{l})$	(I)	Disproportionation reaction
B	$2\text{NaH}(\text{s}) \xrightarrow{\Delta} 2\text{Na}(\text{s}) + \text{H}_2(\text{g})$	(II)	Combination reaction
C	$\text{V}_2\text{O}_5(\text{s}) + 5\text{Ca}(\text{s}) \xrightarrow{\Delta} 2\text{V}(\text{s}) + 5\text{CaO}(\text{s})$	(III)	Decomposition reaction
D	$2\text{H}_2\text{O}_2(\text{aq}) \xrightarrow{\Delta} 2\text{H}_2\text{O}(\text{l}) + \text{O}_2(\text{g})$	(IV)	Displacement reaction

Choose the **correct** answer from the options given below :

- (1) A-II, B-III, C-IV, D-I  
 (2) A-II, B-III, C-I, D-IV  
 (3) A-III, B-IV, C-I, D-II  
 (4) A-IV, B-I, C-II, D-III

Ans. (1)

Sol. (A) Combustion of hydrocarbon


(B) Decomposition into gaseous product.


(C) Displacement of 'V' by 'Ca' atom.


(D) Disproportionation of  $\text{H}_2\text{O}_2^{-1}$  into  $\text{O}^{-2}$  and  $\text{O}^0$  oxidation states.



54. Given below are two statements :

**Statement I :**  will undergo alkaline hydrolysis at a faster rate than




**Statement II :** , intramolecular substitution takes place first by involving lone pair of electrons on nitrogen.

In the light of the above statements, choose the **most appropriate** answer from the options given below :

- (1) Both Statement I and Statement II are incorrect
- (2) Statement I is incorrect but statement II is correct
- (3) Both Statement I and Statement II are correct
- (4) Statement I is correct but Statement II is incorrect

Ans. (3)




Rate of (a) is faster than rate of (b) because it is an intramolecular substitution.

55. A weak acid HA has degree of dissociation  $x$ . Which option gives the correct expression of  $\text{pH} = \text{pK}_a$  ?

- (1)  $\log(1 + 2x)$
- (2)  $\log\left(\frac{1-x}{x}\right)$
- (3) 0
- (4)  $\log\left(\frac{x}{1-x}\right)$

Ans. (4)



$t=0$  a

$t=t$   $a(1-x)$   $ax$   $ax$

$$K_a = (ax) \frac{(x)}{1-x}; [\text{H}^+] = ax$$

$$-\log(K_a) = -\log(ax) - \log\left(\frac{x}{1-x}\right)$$

$$\text{pK}_a = \text{pH} - \log\left(\frac{x}{1-x}\right)$$

$$\text{pH} - \text{pK}_a = \log\left(\frac{x}{1-x}\right)$$

56. Consider 'n' is the number of lone pair of electrons present in the equatorial position of the most stable structure of  $\text{ClF}_3$ . The ions from the following with 'n' number of unpaired electrons are :


- A.  $\text{V}^{3+}$
- B.  $\text{Ti}^{3+}$
- C.  $\text{Cu}^{2+}$
- D.  $\text{Ni}^{2+}$
- E.  $\text{Ti}^{2+}$

Choose the **correct** answer from the options given below :

- (1) A and C only
- (2) A, D and E only
- (3) B and C only
- (4) B and D only

Ans. (2)

**Sol.**  $\text{ClF}_3$



$n = 2$  (No of lone pair present in equatorial plane)  
(Unpaired  $e^-$ )

- (A)  $\text{V}^{3+} : [\text{Ar}]3d^2$  2
- (B)  $\text{Ti}^{3+} : [\text{Ar}]3d^1$  1
- (C)  $\text{Cu}^{2+} : [\text{Ar}]3d^9$  1
- (D)  $\text{Ni}^{2+} : [\text{Ar}]3d^8$  2
- (E)  $\text{Ti}^{2+} : [\text{Ar}]3d^2$  2

57.

$[\text{A}]_0 / \text{molL}^{-1}$	$t_{1/2} / \text{min}$
0.100	200
0.025	100

For a given reaction  $\text{R} \rightarrow \text{P}$ ,  $t_{1/2}$  is related to  $[\text{A}]_0$  as given in table :

Given :  $\log 2 = 0.30$

Which of the following is **true** ?

- A. The order of the reaction is  $\frac{1}{2}$ .
  - B. If  $[\text{A}]_0$  is 1M, then  $t_{1/2}$  is  $200\sqrt{10}$  min
  - C. The order of the reaction changes to 1 if the concentration of reactant changes from 0.100 M to 0.500 M.
  - D.  $t_{1/2}$  is 800 min for  $[\text{A}]_0 = 1.6$  M
- Choose the **correct** answer from the options given below :

- (1) A and C only
- (2) A and B only
- (3) A, B and D only
- (4) C and D only

Ans. (3)



Sol.  $t_{1/2} \propto \frac{1}{A_0^{n-1}}$

$$\frac{(t_{1/2})_1}{(t_{1/2})_2} = \frac{(A_0)_2^{n-1}}{(A_0)_1^{n-1}}$$

$$\frac{200}{100} = \left( \frac{0.025}{0.100} \right)^{n-1}$$

$$2 = \left( \frac{1}{4} \right)^{n-1}$$

$$n - 1 = -\frac{1}{2}$$

$$n = \frac{1}{2} \text{ (order)}$$

$$\Rightarrow t_{1/2} \propto \sqrt{A_0}$$

$$\frac{200}{t_{1/2}} = \frac{(0.1)^{1/2}}{(1)^{1/2}}$$

$$\text{when } A_0 = 1 \text{ M}$$

$$t_{1/2} = 200\sqrt{10} \text{ min}$$


\* 1st order kinetics have  $t_{1/2}$  independent of their concentration. So upon changing the concentration  $t_{1/2}$  should not change for first order reaction.

$$\frac{200}{t_{1/2}} = \frac{(0.1)^{1/2}}{(1.6)^{1/2}}$$





$$\text{when } A_0 = 1.6 \text{ M}$$

$$t_{1/2} = 800 \text{ min}$$

58. A molecule ("P") on treatment with acid undergoes rearrangement and gives ("Q") ("Q") on ozonolysis followed by reflux under alkaline condition gives ("R"). The structure of ("R") is given below :




The structure of ("P") is



Ans. (2)

Sol.




Note : In question about molecule "P" is not clarified, whether it is alcohol or alkene and as in question language rearrangement product is asking hence according to question language ans. is either (2) or (4). As alkene also undergoes rearrangement in presence of acid but option (2) also fulfil all conditions.



59. Ice and water are placed in a closed container at a pressure of 1 atm and temperature 273.15 K. If pressure of the system is increased 2 times, keeping temperature constant, then identify correct observation from following :

- (1) Volume of system increases.
- (2) Liquid phase disappears completely.
- (3) The amount of ice decreases.
- (4) The solid phase (ice) disappears completely.


Ans. (4)



If pressure is made two time then mixture of ice and water will completely convert into water (liquid) form.

60. The molecules having square pyramidal geometry are
- (1)  $BrF_5$  &  $XeOF_4$
  - (2)  $SbF_5$  &  $XeOF_4$
  - (3)  $SbF_5$  &  $PCl_5$
  - (4)  $BrF_5$  &  $PCl_5$

Ans. (1)



Sol.


$BrF_5$  : Square pyramidal  
 $XeOF_4$  : Square pyramidal  
 $SbF_5$  : Trigonal bipyramidal  
 $PCl_5$  : Trigonal bipyramidal

61. The metal ion whose electronic configuration is not affected by the nature of the ligand and which gives a violet colour in non-luminous flame under hot condition in borax bead test is

- (1)  $Ti^{3+}$
- (2)  $Ni^{2+}$
- (3)  $Mn^{2+}$
- (4)  $Cr^{3+}$

Ans. (2)

Sol.  $Ni^{+2}$  gives violet coloured bead in non-luminous flame under hot conditions.  $Ni^{+2}$  has  $d^8$  configuration which does not depend on nature of ligand present in octahedral complex.



62. Both acetaldehyde and acetone (individually) undergo which of the following reactions?

- A. Iodoform Reaction
- B. Cannizaro Reaction
- C. Aldol condensation
- D. Tollen's Test
- E. Clemmensen Reduction

Choose the **correct** answer from the options given below :

- (1) A, B and D only
- (2) A, C and E only
- (3) C and E only
- (4) B, C and D only

Ans. (2)

Sol.

S.No.	Name of Reaction	Acetaldehyde $CH_3-C(=O)-H$	Acetone $CH_3-C(=O)-CH_3$
1	Iodoform reaction	⊕ve	⊕ve
2	Cannizaro	⊖ve	⊖ve
3	Aldol reaction	⊕ve	⊕ve
4	Tollen's test	⊕ve	⊖ve
5	Clemmensen reduction	⊕ve	⊕ve

Ans. (2) A, C and E only



63. In a multielectron atom, which of the following orbitals described by three quantum numbers with have same energy in absence of electric and magnetic fields?

- A.  $n = 1, l = 0, m_l = 0$   
 B.  $n = 2, l = 0, m_l = 0$   
 C.  $n = 2, l = 1, m_l = 1$   
 D.  $n = 3, l = 2, m_l = 1$   
 E.  $n = 3, l = 2, m_l = 0$

Choose the **correct** answer from the options given below :

- (1) A and B only  
 (2) B and C only  
 (3) C and D only  
 (4) D and E only


**Ans. (4)**

**Sol.**

	orbital
A : $n = 1, l = 0, m_l = 0$	1s
B : $n = 2, l = 0, m_l = 0$	2s
C : $n = 3, l = 1, m_l = 1$	3p
D : $n = 3, l = 2, m_l = 1$	3d
E : $n = 3, l = 2, m_l = 0$	3d



In absence of electric and magnetic fields, all orbitals of 3d are degenerate

64. The products A and B in the following reactions, respectively are



- (1)  $\text{CH}_3-\text{CH}_2-\text{CH}_2-\text{ONO}$ ,  $\text{CH}_3-\text{CH}_2-\text{CH}_2-\text{NC}$   
 (2)  $\text{CH}_3-\text{CH}_2-\text{CH}_2-\text{ONO}$ ,  $\text{CH}_3-\text{CH}_2-\text{CH}_2-\text{CN}$   
 (3)  $\text{CH}_3-\text{CH}_2-\text{CH}_2-\text{NO}_2$ ,  $\text{CH}_3-\text{CH}_2-\text{CH}_2-\text{CN}$   
 (4)  $\text{CH}_3-\text{CH}_2-\text{CH}_2-\text{NO}_2$ ,  $\text{CH}_3-\text{CH}_2-\text{CH}_2-\text{NC}$

**Ans. (4)**



65. What is the freezing point depression constant of a solvent, 50 g of which contain 1 g non volatile solute (molar mass  $256 \text{ g mol}^{-1}$ ) and the decrease in freezing point is  $0.40 \text{ K}$ ?

- (1)  $5.12 \text{ K kg mol}^{-1}$  (2)  $4.43 \text{ K kg mol}^{-1}$   
 (3)  $1.86 \text{ K kg mol}^{-1}$  (4)  $3.72 \text{ K kg mol}^{-1}$

**Ans. (1)**

**Sol.**  $\Delta T_f = K_b \cdot m$

$$0.4 = K_b \cdot \frac{1}{50 \times 10^{-3} \times \frac{256}{256}}$$

$$K_b = 5.12 \text{ K kg / mol}$$

66. Consider the following elements In, Tl, Al, Pb, Sn and Ge.

The most stable oxidation states of elements with highest and lowest first ionisation enthalpies, respectively, are


- (1) +2 and +3 (2) +4 and +3  
 (3) +4 and +1 (4) +1 and +4

**Ans. (3)**

**Sol.** Among Al, In, Tl, Ge, Sn, Pb, the metal having highest  $IE_1$  is Ge and lowest  $IE_1$  is In.



Most stable oxidation state of Ge is +4 and In is +3.

67. The correct order of stability of following carbocations is :



- (1)  $A > B > C > D$  (2)  $B > C > A > D$   
 (3)  $C > B > A > D$  (4)  $C > A > B > D$

**Ans. (4)**



Solution :-

C is aromatic due to  $\oplus$ ve charge hence it is most stable

A have more resonance structure

B have less resonance structure






D have only hyper conjugation

Consider First Aromaticity > Resonance > Hyper conjugation

Ans.  $D < B < A < C$



68. The compounds that produce  $\text{CO}_2$  with aqueous  $\text{NaHCO}_3$  solution are :



Choose the **correct** answer from the options given below :

- (1) A and C only                      (2) A, B and E only  
(3) A, C and D only                (4) A and B only

Ans. (3)

Sol. A, C, D produce  $\text{CO}_2$  with aqueous  $\text{NaHCO}_3$  solution.

A, C, D acids are stronger acid than  $\text{H}_2\text{CO}_3$  (Carbonic acid)




69. Which of the following oxidation reactions are carried out by both  $\text{K}_2\text{Cr}_2\text{O}_7$  and  $\text{KMnO}_4$  in acidic medium ?

- A.  $\text{I}^- \rightarrow \text{I}_2$   
B.  $\text{S}^{2-} \rightarrow \text{S}$   
C.  $\text{Fe}^{2+} \rightarrow \text{Fe}^{3+}$   
D.  $\text{I}^- \rightarrow \text{IO}_3^-$   
E.  $\text{S}_2\text{O}_3^{2-} \rightarrow \text{SO}_4^{2-}$

Choose the **correct** answer from the options given below :

- (1) B, C and D only                (2) A, D and E only  
(3) A, B and C only                (4) C, D and E only

Ans. (3)



70. Given below are two statements :

**Statement I :** D-glucose pentaacetate reacts with 2, 4-dinitrophenylhydrazine.

**Statement II :** Starch, on heating with concentrated sulfuric acid at  $100^\circ\text{C}$  and 2-3 atmosphere pressure produces glucose.

In the light of the above statements, choose the **correct** answer from the options given below


- (1) Both Statement I and Statement II are false  
(2) Statement I is false but Statement II is true  
(3) Statement I is true but Statement II is false  
(4) Both Statement I and Statement II are true

Ans. (2)

Sol.

### SECTION-B

71. Given below is the plot of the molar conductivity vs  $\sqrt{\text{concentration}}$  for KCl in aqueous solution.



If, for the higher concentration of KCl solution, the resistance of the conductivity cell is  $100\Omega$ , then the resistance of the same cell with the dilute solution is ' $x$ '  $\Omega$ .

The value of  $x$  is \_\_\_\_\_ (Nearest integer)

Ans. 150



**Sol.**  $R = \rho \frac{\ell}{A}$

$$\kappa = G \cdot G^* \quad G = \frac{1}{R}; \kappa = \frac{1}{\rho}$$

$$G^* = \frac{\ell}{A}$$

R = Resistance

$\rho$  = Resistivity

$$\frac{\ell}{A} = \text{cell constant } (G^*)$$

$$\frac{\kappa_c}{\kappa_d} = \frac{R_d}{R_c}; \lambda_m = \frac{\kappa \times 1000}{C}$$

$$\frac{\kappa_c}{\kappa_d} = \frac{(\lambda_m \cdot C)}{(\lambda_m \cdot C)_d} = \frac{R_d}{R_c} \quad \begin{array}{l} c = \text{concentrated sol.} \\ d = \text{diluted solution} \end{array}$$

$$\frac{100 \cdot (0.15)^2}{150 \cdot (0.1)^2} = \frac{R_d}{100}$$

$$R_d = 150 \Omega$$

- 72.** Quantitative analysis of an organic compound (X) shows following % composition.

C : 14.5%                      Cl : 64.46%

H : 1.8%

(Empirical formula mass of the compound (X) is  $\times 10^{-1}$ )

(Given molar mass in  $\text{g mol}^{-1}$  of C : 12, H : 1, O : 16, Cl : 35.5)

**Ans. 1655**

**Sol.**

	C	Cl	H	O
%mass	14.5	64.46	1.8	19.24
Molar ratio	$\frac{14.5}{12}$	$\frac{64.46}{35.5}$	$\frac{1.8}{1}$	$\frac{19.24}{16}$
	1.2	1.8	1.8	1.2
Minimum	2	3	3	2
integral ratio				

Empirical formula =  $\text{C}_2\text{H}_3\text{Cl}_3\text{O}_2$

Mass = 165.5

Mass =  $1655 \times 10^{-1}$

- 73.** The molarity of a 70% (mass/mass) aqueous solution of a monobasic acid (X) is \_\_\_\_\_ M (Nearest integer)

[Given : Density of aqueous solution of (X) is  $1.25 \text{ g mL}^{-1}$ ]

Molar mass of the acid is  $70 \text{ g mol}^{-1}$ ]


**Ans. 125**

**Sol.** Assuming 100 gm solution contain 70 gm solute.

Volume of 100 gm solution will be  $\frac{100}{1.25}$  ml.

$$\text{Molarity} = \frac{70/70}{100/1.25} \times 1000 = 12.5 \text{ or } 125 \times 10^{-1}$$

- 74.** Consider the following sequence of reactions :




Chlorobenzene

11.25 mg of chlorobenzene will produce  $\times 10^{-1}$  mg of product B.

(Consider the reactions result in complete conversion.)

[Given molar mass of C, H, O, N and Cl as 12, 1, 16, 14 and  $35.5 \text{ g mol}^{-1}$  respectively]

**Ans. 93**





$$\frac{11.25 \times 10^{-3}}{112.5} = \frac{x \times 10^{-1} \times 10^{-3}}{93}$$

$$x \times 10^{-1} = 93 \times 0.1$$

$$x = 93 \text{ mg}$$

- 75.** The formation enthalpies,  $\Delta H_f^\ominus$  for  $\text{H}_{(\text{g})}$  and  $\text{O}_{(\text{g})}$  are 220.0 and 250.0  $\text{kJ mol}^{-1}$ , respectively, at 298.15 K, and  $\Delta H_f^-$  for  $\text{H}_2\text{O}_{(\text{g})}$  is  $-242.0 \text{ kJ mol}^{-1}$  at the same temperature. The average bond enthalpy of the O–H bond in water at 298.15 K is \_\_\_\_\_  $\text{kJ mol}^{-1}$  (nearest integer).

**Ans. 466**



$$\Delta H_f(\text{H}_2\text{O}_{(\text{l})}) = -242 = 440 + 250 - 2(\text{B.E.}(\text{O}-\text{H}))$$

$$\boxed{\text{BE}(\text{O}-\text{H}) = 466 \text{ KJ/mol}}$$